




## **ROBO4 Monoclonal Antibody**

Catalog No	BYmab-07045
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	ROBO4 UNQ421/PRO3674
Protein Name	Roundabout homolog 4 (Magic roundabout)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ROBO4 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 110kD
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  110kD  integral component of membrane,extracellular exosome,  Specifically expressed in endothelial cells. Expressed in endothelial and intimal
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  110kD  integral component of membrane,extracellular exosome,  Specifically expressed in endothelial cells. Expressed in endothelial and intimal cells of the ascending aorta (PubMed:30455415).  function:Receptor for Slit proteins, at least for SLIT2, and seems to be involved in angiogenesis and vascular patterning. May mediate the inhibition of primary endothelial cell migration by Slit proteins.,similarity:Belongs to the immunoglobulin superfamily. ROBO family.,similarity:Contains 2 fibronectin type-III domains.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Interacts with SLIT2 and ENAH.,tissue specificity:Specifically expressed in endothelial cells. Expressed at sites of angiogenesis in different

Nanjing BYabscience technology Co.,Ltd







	superfamily. ROBO family.,similarity:Contains 2 fibronectin type-III domains.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Interacts with SLIT2 and ENAH.,tissue specificity:Specifically expressed in endothelial cells. Expressed at sites of angiogenesis in different tumor types.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658