



HES1 Monoclonal Antibody

IgG Human;Rat;Mouse WB HES1 BHLHB39 HL HRY Transcription factor HES-1 (Class B basic helix-loop-helix protein 39) (bHLHb39) (Hairy and enhancer of split 1) (Hairy homolog) (Hairy-like protein) (hHL) Synthesized peptide derived from part region of human protein
Human;Rat;Mouse WB HES1 BHLHB39 HL HRY Transcription factor HES-1 (Class B basic helix-loop-helix protein 39) (bHLHb39) (Hairy and enhancer of split 1) (Hairy homolog) (Hairy-like protein) (hHL) Synthesized peptide derived from part region of human protein
WB HES1 BHLHB39 HL HRY Transcription factor HES-1 (Class B basic helix-loop-helix protein 39) (bHLHb39) (Hairy and enhancer of split 1) (Hairy homolog) (Hairy-like protein) (hHL) Synthesized peptide derived from part region of human protein
HES1 BHLHB39 HL HRY Transcription factor HES-1 (Class B basic helix-loop-helix protein 39) (bHLHb39) (Hairy and enhancer of split 1) (Hairy homolog) (Hairy-like protein) (hHL) Synthesized peptide derived from part region of human protein
Transcription factor HES-1 (Class B basic helix-loop-helix protein 39) (bHLHb39) (Hairy and enhancer of split 1) (Hairy homolog) (Hairy-like protein) (hHL) Synthesized peptide derived from part region of human protein
(Hairy and enhancer of split 1) (Hairy homolog) (Hairy-like protein) (hHL) Synthesized peptide derived from part region of human protein
LICA Managanal Antibady detects and grange levels of protein
HES1 Monoclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
30kD
Nucleus.
Kidney,Prostate,Skin,
domain:Has a particular type of basic domain (presence of a helix-interrupting proline) that binds to the N-box (CACNAG), rather than the canonical E-box (CANNTG).,domain:The bHLH, as well as cooperation between the central Orange domain and the C-terminal WRPW motif, is required for transcriptional repressor activity.,domain:The C-terminal WRPW motif is a transcriptional repression domain necessary for the interaction with Groucho/TLE family members, transcriptional corepressors recruited to specific target DNA by Hairy-related proteins.,function:Transcriptional repressor of genes that require a bHLH protein for their transcription. May act as a negative regulator of myogenesis by inhibiting the functions of MYOD1 and ASH1. Binds DNA on N-box motifs: 5'-CACNAG-3' with high affinity and on E-box motifs: 5'-CANNTG-3 with low affinity.,similarity:Contains 1 basic helix-loop-helix (bHLH) d

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Background	This protein belongs to the basic helix-loop-helix family of transcription factors. It is a transcriptional repressor of genes that require a bHLH protein for their transcription. The protein has a particular type of basic domain that contains a helix interrupting protein that binds to the N-box rather than the canonical E-box. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658