



LUM Monoclonal Antibody

Catalog No	BYmab-07096
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	LUM LDC SLRR2D
Protein Name	Lumican (Keratan sulfate proteoglycan lumican) (KSPG lumican)
Immunogen	Synthesized peptide derived from human protein . at AA range: 30-110
Specificity	LUM Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year 37kD
Synonyms	
Synonyms Observed Band	37kD
Synonyms Observed Band Cell Pathway	37kD Secreted, extracellular space, extracellular matrix. Cornea and other tissues. developmental stage:Present in the extracellular matrix of human articular cartilage at all ages, although its abundance is far greater in the adult. In the adult cartilage lumican exists predominantly in a glycoprotein form lacking keratan sulfate, whereas the juvenile form of the molecule is a proteoglycan.,PTM:Sulfated on tyrosine residue(s).,similarity:Belongs to the small leucine-rich proteoglycan (SLRP) family. Class II subfamily.,similarity:Contains 12
Synonyms Observed Band Cell Pathway Tissue Specificity	37kD Secreted, extracellular space, extracellular matrix. Cornea and other tissues. developmental stage:Present in the extracellular matrix of human articular cartilage at all ages, although its abundance is far greater in the adult. In the adult cartilage lumican exists predominantly in a glycoprotein form lacking keratan sulfate, whereas the juvenile form of the molecule is a proteoglycan.,PTM:Sulfated on tyrosine residue(s).,similarity:Belongs to the small leucine-rich proteoglycan (SLRP) family. Class II subfamily.,similarity:Contains 12 LRR (leucine-rich) repeats.,subunit:Binds to laminin.,tissue specificity:Cornea and

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	fibrils and the highly charged hydrophilic glycosaminoglycans regulate interfibrillar spacings. Lumican is the major keratan sulfate proteoglycan of the cornea but is also distributed in interstitial collagenous matrices throughout the body. Lumican may regulate collagen fibril organization and circumferential growth, corneal transparency, and epithelial cell migration and tissue repair. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using LUM Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658