



DBLOH Monoclonal Antibody

| | |
|---------------------------|---|
| Catalog No | BYmab-07205 |
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | DIABLO SMAC |
| Protein Name | Diablo homolog, mitochondrial (Direct IAP-binding protein with low pI) (Second mitochondria-derived activator of caspase) (Smac) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | DBLOH Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 26kD |
| Cell Pathway | Mitochondrion . Released into the cytosol when cells undergo apoptosis. |
| Tissue Specificity | Ubiquitously expressed with highest expression in testis. Expression is also high in heart, liver, kidney, spleen, prostate and ovary. Low in brain, lung, thymus and peripheral blood leukocytes. Isoform 3 is ubiquitously expressed. |
| Function | domain:The mature N-terminus mediates interaction with XIAP.,function:Promotes apoptosis by activating caspases in the cytochrome c/Apaf-1/caspase-9 pathway. Acts by opposing the inhibitory activity of inhibitor of apoptosis proteins (IAP).,subcellular location:But released into the cytosol when cells undergo apoptosis.,subunit:Interacts with NGFRAP1/BEX3 (By similarity). Homodimer. Interacts with BIRC2, BIRC3, XIAP and BIRC7.,tissue specificity:Ubiquitously expressed with highest expression in testis. Expression is also high in heart, liver, kidney, spleen, prostate and ovary. Low in brain, lung, thymus and peripheral blood leukocytes., |

Nanjing BYabscience technology Co.,Ltd



Background

This gene encodes an inhibitor of apoptosis protein (IAP)-binding protein. The encoded mitochondrial protein enters the cytosol when cells undergo apoptosis, and allows activation of caspases by binding to inhibitor of apoptosis proteins. Overexpression of the encoded protein sensitizes tumor cells to apoptosis. A mutation in this gene is associated with young-adult onset of nonsyndromic deafness-64. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, May 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using DBLOH Monoclonal Antibody