



POSTN Monoclonal Antibody

Catalog No	BYmab-07222
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	POSTN OSF2
Protein Name	Periostin (PN) (Osteoblast-specific factor 2) (OSF-2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	POSTN Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	91kD
Cell Pathway	Golgi apparatus . Secreted . Secreted, extracellular space, extracellular matrix . Colocalizes with BMP1 in the Golgi. .
Tissue Specificity	Widely expressed with highest levels in aorta, stomach, lower gastrointestinal tract, placenta, uterus, thyroid tissue and breast. Up-regulated in epithelial ovarian tumors. Not expressed in normal ovaries. Also highly expressed at the tumor periphery of lung carcinoma tissue but not within the tumor. Overexpressed in breast cancers.
Function	function: Binds to heparin. Induces cell attachment and spreading and plays a role in cell adhesion. May play a role in extracellular matrix mineralization. .PTM: Gamma-carboxyglutamate residues are formed by vitamin K dependent carboxylation. These residues are essential for the binding of calcium. .similarity: Contains 1 EMI domain. .similarity: Contains 4 FAS1 domains. .tissue specificity: Widely expressed with highest levels in aorta, stomach, lower gastrointestinal tract, placenta, uterus and breast. Up-regulated in epithelial ovarian tumors. Not expressed in normal ovaries. Also highly expressed

Nanjing BYabscience technology Co.,Ltd



at the tumor periphery of lung carcinoma tissue but not within the tumor.
Overexpressed in breast cancers.,

Background

This gene encodes a secreted extracellular matrix protein that functions in tissue development and regeneration, including wound healing, and ventricular remodeling following myocardial infarction. The encoded protein binds to integrins to support adhesion and migration of epithelial cells. This protein plays a role in cancer stem cell maintenance and metastasis. Mice lacking this gene exhibit cardiac valve disease, and skeletal and dental defects. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images