



PRLR Monoclonal Antibody

Catalog No	BYmab-07228
Isotype	lgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	PRLR
Protein Name	Prolactin receptor (PRL-R)
Immunogen	Synthesized peptide derived from human protein . at AA range: 80-160
Specificity	PRLR Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	68kD
Cell Pathway	Membrane ; Single-pass type I membrane protein .; [Isoform 7]: Secreted.
Tissue Specificity	Expressed in breast, placenta, kidney, liver and pancreas.
Function	domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:This is a receptor for the anterior pituitary hormone prolactin (PRL). Isoforms 4 and 6 are unable to transduce prolactin signaling.,similarity:Belongs to the type I cytokine receptor family. Type 1 subfamily.,similarity:Contains 2 fibronectin type-III domains.,subcellular location:Isoform 7.,subunit:Interacts with SMARCA1. Interacts with GH1.,tissue specificity:Expressed in breast, placenta, kidney, liver and pancreas.,
Background	This gene encodes a receptor for the anterior pituitary hormone, prolactin, and belongs to the type I cytokine receptor family. Prolactin-dependent signaling
	Naniing BYabscience technology Co. 1td

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务,欢迎咨询

	occurs as the result of ligand-induced dimerization of the prolactin receptor. Several alternatively spliced transcript variants encoding different membrane-bound and soluble isoforms have been described for this gene, which may function to modulate the endocrine and autocrine effects of prolactin in normal tissue and cancer. [provided by RefSeq, Feb 2011],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
Products Images		

Nanjing BYabscience technology Co.,Ltd