



SPRE1 Monoclonal Antibody

Catalog No	BYmab-07258
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	SPRED1
Protein Name	Sprouty-related, EVH1 domain-containing protein 1 (Spred-1) (hSpred1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 210-290
Specificity	SPRE1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	48kD
Cell Pathway	Cell membrane ; Peripheral membrane protein . Membrane, caveola ; Peripheral membrane protein . Nucleus . Localized in cholesterol-rich membrane raft/caveola fractions.
Tissue Specificity	Weakly expressed in embryonic cell line HEK293.
Function	disease:Defects in SPRED1 are the cause of neurofibromatosis type 1-like syndrome (NFLS) [MIM:611431]. Neurofibromatosis type 1 (NF1) is one of the most frequent autosomal dominant diseases. It belongs to the group of disorders known as the 'neuro-cardio-facial-cutaneous' syndromes, present with a variable degree of cognitive impairment, facial dysmorphism, congenital heart defects and skin abnormalities. NFLS is a form of these disorders with autosomal dominant trait consisting of multiple cafe-au-lait spots, axillary freckling, macrocephaly and a Noonan-like dysmorphism in some individuals.,function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase. Negatively regulates hematopoiesis of bone marrow.,PTM:Phosphorylated on tyrosine.,sequence caution:Contaminating sequence. Potential poly-A

Nanjing BYabscience technology Co.,Ltd



sequence.,similarity:Contains 1 KBD domain.,similarity:Contains

Background

The protein encoded by this gene is a member of the Sprouty family of proteins and is phosphorylated by tyrosine kinase in response to several growth factors. The encoded protein can act as a homodimer or as a heterodimer with SPRED2 to regulate activation of the MAP kinase cascade. Defects in this gene are a cause of neurofibromatosis type 1-like syndrome (NFLS). [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658