




SIAL Monoclonal Antibody

| Catalog No | BYmab-07266 |
|--------------------|--|
| Isotype | lgG |
| Reactivity | Human;Rat;Mouse |
| Applications | WB |
| Gene Name | IBSP BNSP |
| Protein Name | Bone sialoprotein 2 (Bone sialoprotein II) (BSP II) (Cell-binding sialoprotein) (Integrin-binding sialoprotein) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 230-310 |
| Specificity | SIAL Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 34kD |
| Cell Pathway | Secreted. |
| Tissue Specificity | |
| Function | function:Binds tightly to hydroxyapatite. Appears to form an integral part of the mineralized matrix. Probably important to cell-matrix interaction. Promotes Arg-Gly-Asp-dependent cell attachment.,miscellaneous:It is possible that the segments of clustered carboxyl groups mediate the strong binding to hydroxyapatite.,PTM:N-glycosylated; glycans consist of sialated and core-fucosylated bi-, tri- and tetraantennary chains.,PTM:O-glycosylated at eight sites; mucin-type glycans contain Gal, GlcNAc, GalNAc and terminal NeuAc.,PTM:Sulfated on either Tyr-313 or Tyr-314., |
| Background | The protein encoded by this gene is a major structural protein of the bone matrix. It constitutes approximately 12% of the noncollagenous proteins in human bone and is synthesized by skeletal-associated cell types, including hypertrophic |
| | Nanjing BYabscience technology Co.,Ltd |



Products Images



Nanjing BYabscience technology Co.,Ltd