



PLAP Monoclonal Antibody

IsotypeIgGReactivityHuman;Mouse;RatApplicationsWBGene NamePLAA PLAPProtein NamePhospholipase A-2-activating protein (PLA2P) (PLAP)ImmunogenSynthesized peptide derived from human protein . at AA range: 531-580SpecificityPLAP Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%
ApplicationsWBGene NamePLAA PLAPProtein NamePhospholipase A-2-activating protein (PLA2P) (PLAP)ImmunogenSynthesized peptide derived from human protein . at AA range: 531-580SpecificityPLAP Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000
Gene NamePLAA PLAPProtein NamePhospholipase A-2-activating protein (PLA2P) (PLAP)ImmunogenSynthesized peptide derived from human protein . at AA range: 531-580SpecificityPLAP Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/ml
Protein NamePhospholipase A-2-activating protein (PLA2P) (PLAP)ImmunogenSynthesized peptide derived from human protein . at AA range: 531-580SpecificityPLAP Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/ml
ImmunogenSynthesized peptide derived from human protein . at AA range: 531-580SpecificityPLAP Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/ml
SpecificityPLAP Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/ml
FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/ml
SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/ml
PurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/ml
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml
Concentration 1 mg/ml
Purity ≥90%
Storage Stability -20°C/1 year
Synonyms
Observed Band 87kD
Cell Pathway Nucleus . Cytoplasm . Cell junction, synapse . Recruited to damaged lysoson decorated with K48-linked ubiquitin chains
Tissue Specificity Fetal brain,Kidney,Testis,Uterus,
Function function:Plays an important role in the regulation of specific inflammatory dise processes.,sequence caution:Translated as Gln.,similarity:Belongs to the WD repeat PLAP family.,similarity:Contains 1 PFU domain.,similarity:Contains 1 F domain.,similarity:Contains 7 WD repeats.,
Background function:Plays an important role in the regulation of specific inflammatory dise processes.,sequence caution:Translated as Gln.,similarity:Belongs to the WD repeat PLAP family.,similarity:Contains 1 PFU domain.,similarity:Contains 1 F domain.,similarity:Contains 7 WD repeats.,
matters needing Avoid repeated freezing and thawing! attention Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd