


SPXN Monoclonal Antibody

Catalog No	BYmab-07329
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	C12orf39
Protein Name	Spexin (NPQ)
Immunogen	Synthesized peptide derived from human protein . at AA range: 1-50
Specificity	SPXN Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Observed Band	12kD
Cell Pathway	Secreted. Secreted, extracellular space. Cytoplasmic vesicle, secretory vesicle. Secreted via the classical ER/Golgi-dependent pathway into the extracellular medium largely as a full-length protein without the signal peptide, and not as a hydrolyzed and amidated peptide (PubMed:19193193 and PubMed:17284679). Localized extracellularly surrounding the villous trophoblastic cells. Detected in the serum.
	Secreted. Secreted, extracellular space. Cytoplasmic vesicle, secretory vesicle. Secreted via the classical ER/Golgi-dependent pathway into the extracellular medium largely as a full-length protein without the signal peptide, and not as a hydrolyzed and amidated peptide (PubMed:19193193 and PubMed:17284679). Localized extracellularly surrounding the villous trophoblastic cells. Detected in
Cell Pathway	Secreted. Secreted, extracellular space. Cytoplasmic vesicle, secretory vesicle. Secreted via the classical ER/Golgi-dependent pathway into the extracellular medium largely as a full-length protein without the signal peptide, and not as a hydrolyzed and amidated peptide (PubMed:19193193 and PubMed:17284679). Localized extracellularly surrounding the villous trophoblastic cells. Detected in the serum. Expressed in the type I glomic cells within the carotid body (at protein level). Expressed predominantly in pancreas, testis, kidney, brain and placenta.
Cell Pathway Tissue Specificity	Secreted. Secreted, extracellular space. Cytoplasmic vesicle, secretory vesicle. Secreted via the classical ER/Golgi-dependent pathway into the extracellular medium largely as a full-length protein without the signal peptide, and not as a hydrolyzed and amidated peptide (PubMed:19193193 and PubMed:17284679). Localized extracellularly surrounding the villous trophoblastic cells. Detected in the serum. Expressed in the type I glomic cells within the carotid body (at protein level). Expressed predominantly in pancreas, testis, kidney, brain and placenta. Expressed in submucosal layer of esophagus and stomach fundus. function:Induces contraction of stomach muscle.,tissue specificity:Expressed in submucosal layer of esophagus and stomach fundus. Expressed in brain,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







RefSeq, Feb 2016],

matters	needing	
attention		

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

