



## **ZCCHV Monoclonal Antibody**

Catalog No	BYmab-07355
lsotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	ZC3HAV1 ZC3HDC2 PRO1677
Protein Name	Zinc finger CCCH-type antiviral protein 1 (ADP-ribosyltransferase diphtheria toxin-like 13) (ARTD13) (Zinc finger CCCH domain-containing protein 2) (Zinc finger antiviral protein) (ZAP)
Immunogen	Synthesized peptide derived from human protein . at AA range: 211-260
Specificity	ZCCHV Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	99kD
Cell Pathway	[Isoform 1]: Cytoplasm . Nucleus . Localizes in the cytoplasm at steady state, but shuttles between nucleus and cytoplasm in a XPO1-dependent manner; [Isoform 2]: Cytoplasm .
Tissue Specificity	Brain,Endometrium,Epithelium,Fetal liver,Kidney,Ovarian car
Function	domain:The second and fourth zinc fingers are involved in binding to specific viral RNAs.,function:Induces an innate immunity to viral infections by preventing the accumulation of viral RNAs in the cytoplasm. Seems to recruit the RNA processing exosome to degrade the target RNAs. Inhibits alphavirus and filovirus replication.,similarity:Contains 1 PARP catalytic domain.,similarity:Contains 1 WWE domain.,similarity:Contains 4 C3H1-type zinc fingers.,subcellular location:Localizes in the cytoplasm at steady state, but shuttles between nucleus and cytoplasm in a XPO1-dependent manner.,subunit:Interacts with EXOSC5.,

## Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务,欢迎咨询



Background matters needing	This gene encodes a CCCH-type zinc finger protein that is thought to prevent infection by retroviruses. Studies of the rat homolog indicate that the protein may primarily function to inhibit viral gene expression and induce an innate immunity to viral infection. Alternative splicing occurs at this locus and two variants, each encoding distinct isoforms, are described. [provided by RefSeq, Jul 2008], Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
Products Images	

Nanjing BYabscience technology Co.,Ltd