



GPER Monoclonal Antibody

Catalog No	BYmab-07379
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GPER CEPR CMKRL2 DRY12 GPR30
Protein Name	G-protein coupled estrogen receptor 1 (Chemoattractant receptor-like 2) (Flow-induced endothelial G-protein coupled receptor 1) (FEG-1) (G-protein coupled receptor 30) (GPCR-BR) (IL8-related receptor)
Immunogen	Synthesized peptide derived from human protein . at AA range: 300-380
Specificity	GPER Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	41kD
Cell Pathway	Nucleus. Cytoplasm . Cytoplasm, perinuclear region. Cytoplasm, cytoskeleton. Cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane; Multi-pass membrane protein. Early endosome. Recycling endosome. Golgi apparatus membrane ; Multi-pass membrane protein . Golgi apparatus, trans-Golgi network. Endoplasmic reticulum membrane ; Multi-pass membrane protein. Cell projection, dendrite . Cell projection, dendritic spine membrane ; Multi-pass membrane protein . Cell projection, axon . Cell junction, synapse, postsynaptic density . Mitochondrion membrane ; Multi-pass membrane protein . Colocalized with BSN to the active zone of presynaptic density. Colocalized with DLG4/PSD95 and neurabin-2 PPP1R9B in neuronal synaptosomes (By
Tissue Specificity	Expressed in placenta, endothelial and epithelial cells, non laboring and laboring term myometrium, fibroblasts and cancer-associated fibroblasts (CAF), prostate cancer cells and invasive adenocarcinoma (at protein level). Ubiquitously expressed, but is most abundant in placenta. In brain regions, expressed as a 2.8

Nanjing BYabscience technology Co.,Ltd



kb transcript in basal forebrain, frontal cortex, thalamus, hippocampus, caudate and putamen.

Function

function: Receptor for estrogen., similarity: Belongs to the G-protein coupled receptor 1 family., subcellular location: Protein has been detected in the cell membrane, endoplasmic reticulum and Golgi apparatus. It is currently unclear whether this is a cell surface or intracellular receptor., tissue specificity: Ubiquitously expressed, but is most abundant in placenta. In brain regions, expressed as a 2.8 kb transcript in basal forebrain, frontal cortex, thalamus, hippocampus, caudate and putamen.,

Background

This gene is a member of the G-protein coupled receptor 1 family and encodes a multi-pass membrane protein that localizes to the endoplasmic reticulum. The protein binds estrogen, resulting in intracellular calcium mobilization and synthesis of phosphatidylinositol 3,4,5-trisphosphate in the nucleus. This protein therefore plays a role in the rapid nongenomic signaling events widely observed following stimulation of cells and tissues with estrogen. Alternate transcriptional splice variants which encode the same protein have been characterized. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GPER Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd