



## OR4X2 Monoclonal Antibody

protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene., similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated		
Reactivity Human;Rat;Mouse;  Applications WB  Gene Name OR4X2  Protein Name Olfactory receptor 4X2 (Olfactory receptor OR11-105)  Immunogen Synthesized peptide derived from human protein . at AA range: 220-300  Specificity OR4X2 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptorpolymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene., similarity. Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors of a darge family of G-protein-coupled receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors of a darge family of G-protein-coupled receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors of a darge family of G-protein-coupled receptor share a 7-transmembrane domain structure with many neurotr	Catalog No	BYmab-07492
Applications WB Gene Name OR4X2 Protein Name Olfactory receptor 4X2 (Olfactory receptor OR11-105) Immunogen Synthesized peptide derived from human protein . at AA range: 220-300 Specificity OR4X2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 33kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor _polymorphism: A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene, similarity. Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors Share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain st	Isotype	IgG
Gene Name OR4X2  Protein Name Olfactory receptor 4X2 (Olfactory receptor OR11-105)  Immunogen Synthesized peptide derived from human protein . at AA range: 220-300  Specificity OR4X2 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function: Odorant receptorpolymorphism: A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene, similarity. Belongs to the G-protein coupled receptor 1 family.  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 4X2 (Olfactory receptor OR11-105)  Immunogen Synthesized peptide derived from human protein . at AA range: 220-300  Specificity OR4X2 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptorpolymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene, similarity. Belongs to the G-protein coupled receptor 1 family .,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a 7-transmembrane domain structure with many neuroriansmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of dodrant signals. The olfactory receptor gene family is the largest in	Applications	WB
Immunogen         Synthesized peptide derived from human protein . at AA range: 220-300           Specificity         OR4X2 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Observed Band         33kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Function         function:Odorant receptor "polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene, similarly-Belongs to the G-protein-coupled receptor 1 family.           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor folactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of dodrant signals. The olfactory receptor gene family is the larg	Gene Name	OR4X2
Specificity OR4X2 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptorpolymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogenesimilarity:Belongs to the G-protein coupled receptor 1 family  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Protein Name	Olfactory receptor 4X2 (Olfactory receptor OR11-105)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptorpolymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogenesimilarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptor (GPCR) arising from single coding-exon genes. Olfactory receptores share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Immunogen	Synthesized peptide derived from human protein . at AA range: 220-300
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Cell Pathway           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         function:Odorant receptorpolymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (harmone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Specificity	OR4X2 Monoclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  33kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptorpolymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene., similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor "polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cell membrane; Multi-pass membrane protein.         Tissue Specificity         Function       function:Odorant receptor,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,         Background       Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptorpolymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Dilution	WB 1:500-2000
Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Concentration	1 mg/ml
Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Purity	≥90%
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Storage Stability	-20°C/1 year
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Synonyms	
Function  function:Odorant receptor .,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Observed Band	33kD
Function  function:Odorant receptor .,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Cell Pathway	Cell membrane; Multi-pass membrane protein.
protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene., similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Tissue Specificity	
neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in	Function	function:Odorant receptor .,polymorphism:A stop codon in the gene coding for this protein at position Tyr-27 is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,
	Background	neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using OR4X2 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658