



OR6X1 Monoclonal Antibody

Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated		
Reactivity Human;Rat;Mouse; Applications WB Gene Name OR6X1 Protein Name Olfactory receptor 6X1 (Olfactory receptor OR11-270) Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity OR6X1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a remembers of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a remembers of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a remembers of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a remembers of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors efamily is the largest	Catalog No	BYmab-07515
Applications WB Gene Name OR6X1 Protein Name Olfactory receptor 6X1 (Olfactory receptor OR11-270) Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity OR6X1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are memblers of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors efamily is the largest	Isotype	IgG
Gene Name OR6X1 Protein Name Olfactory receptor 6X1 (Olfactory receptor OR11-270) Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity OR6X1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family. Background Olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptor share a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of g-protein-coupled receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduct	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 6X1 (Olfactory receptor OR11-270) Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity OR6X1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) is the largest transduction of odorant signals. The olfactory receptor gene family is the largest	Applications	WB
Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity OR6X1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family. Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to intilate a neuronal response that tringers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors gene family is the largest transduction of odorant signals. The olfactory gene family is the largest	Gene Name	OR6X1
Specificity OR6X1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Protein Name	Olfactory receptor 6X1 (Olfactory receptor OR11-270)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The offactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory gene family is the largest	Immunogen	Synthesized peptide derived from human protein . at AA range: 230-310
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Specificity	OR6X1 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor yarising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Source	Monoclonal, Mouse,lgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Dilution	WB 1:500-2000
Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Concentration	1 mg/ml
Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Purity	≥90%
Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Synonyms	
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Observed Band	34kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background olfactory receptor family 6 subfamily X member 1(OR6X1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Tissue Specificity	
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest	Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
	Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in

Nanjing BYabscience technology Co.,Ltd





	proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658