



P2Y12 Monoclonal Antibody

Catalog No	BYmab-07533
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	P2RY12 HORK3
Protein Name	P2Y purinoceptor 12 (P2Y12) (ADP-glucose receptor) (ADPG-R) (P2T(AC)) (P2Y(AC)) (P2Y(cyc)) (P2Y12 platelet ADP receptor) (P2Y(ADP)) (SP1999)
Immunogen	Synthesized peptide derived from human protein . at AA range: 110-190
Specificity	P2Y12 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	37kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Highly expressed in the platelets, lower levels in the brain. Lowest levels in the lung, appendix, pituitary and adrenal gland. Expressed in the spinal cord and in the fetal brain.
Function	disease:Defects in P2RY12 are a cause of bleeding disorder [MIM:609821].,function:Receptor for ADP and ATP coupled to G-proteins that inhibit the adenylyl cyclase second messenger system. Not activated by UDP and UTP. Involved in platelets aggregation.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Highly expressed in the platelets, lower levels in the brain. Lowest levels in the lung, appendix, pituitary and adrenal gland. Expressed in the spinal cord and in the fetal brain.,
Background	The product of this gene belongs to the family of G-protein coupled receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine

Nanjing BYabscience technology Co.,Ltd



nucleotides. This receptor is involved in platelet aggregation, and is a potential target for the treatment of thromboembolisms and other clotting disorders. Mutations in this gene are implicated in bleeding disorder, platelet type 8 (BDPLT8). Alternative splicing results in multiple transcript variants of this gene. [provided by RefSeq, Jul 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using P2Y12 Monoclonal Antibody