



O52B4 Monoclonal Antibody

gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in		
ReactivityHuman;Rat;Mouse;ApplicationsWBGene NameOR52B4Protein NameOlfactory receptor 52B4 (Olfactory receptor OR11-3)ImmunogenSynthesized peptide derived from human protein . at AA range: 30-110SpecificityO52B4 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurify≥90%Storage Stability-20°C/1 yearSynonymsCell membrane; Multi-pass membrane protein.Tissue SpecificityCell membrane; Multi-pass membrane protein.Functionfunction:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family.BackgroundOlfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that friggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exol genes. Olfactory receptors proteins and are responsible for the recognition and G protein-mediated	Catalog No	BYmab-07578
Applications WB Gene Name OR52B4 Protein Name Olfactory receptor 52B4 (Olfactory receptor OR11-3) Immunogen Synthesized peptide derived from human protein . at AA range: 30-110 Specificity O52B4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell membrane; Multi-pass membrane protein. Tissue Specificity Cell membrane; Multi-pass membrane protein. Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response the for the recognition and G protein-mediated	Isotype	lgG
Gene Name OR52B4 Protein Name Olfactory receptor 52B4 (Olfactory receptor OR11-3) Immunogen Synthesized peptide derived from human protein . at AA range: 30-110 Specificity O52B4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell membrane; Multi-pass membrane protein. Tissue Specificity Function: Odorant receptorpolymorphism: A stop codon at position Ala-40 in the gene coding for this protein is nere frequent in non-Africans than in African-Americans, similarity: Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptor sinteract with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smill. The offactory receptors subrane domain structure with many neurotansmitter and hormone receptors and are responsible for the recoonfloor of a smill. The offactory receptors indered for the receptor sinteract with many neurotansmitter and hormone receptors and are responsible for the receptor sind are	Reactivity	Human;Rat;Mouse;
Protein NameOlfactory receptor 52B4 (Olfactory receptor OR11-3)ImmunogenSynthesized peptide derived from human protein . at AA range: 30-110SpecificityO52B4 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCell membrane; Multi-pass membrane protein.Charlen SpecificityCell membrane; Multi-pass membrane protein.Functionfunction:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon at molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-cong genes.BackgroundOlfactory receptors interact with odorant molecules in the nose, to infitate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-cong genes. Olfactory receptor share a <i>r</i> -transmembrane domain structure with many neuroransmitter and hormone receptors and are responsible for the recognition and G protein-oupled receptor (GPCR) arising from single coding-cong genes. Olfactory receptors share a <i>r</i> -transmembrane domain structure with many neuroransmitter and hormone receptors and are responsible for the recognition and G protein areadiate	Applications	WB
ImmunogenSynthesized peptide derived from human protein . at AA range: 30-110SpecificityO52B4 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearObserved Band34kDCell PathwayCell membrane; Multi-pass membrane protein.Tissue Specificityfunction: Odorant receptorpolymorphism: A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing paseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs the G-protein coupled receptor 1 family.BackgroundOffactory receptors interact with dorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The offactory receptor arising from single coding exong genes. Offactory receptor share a origenes are members of a large family of G-protein-coupled receptor 1 family.BackgroundOffactory receptors interact with dorant melocules in the nose, to initiate a neuronal response that triggers the perception of a smell. The offactory receptor genes or genes. Offactory receptor share a origens are members of a large family of G-protein-coupled receptor 1 family.	Gene Name	OR52B4
Specificity O52B4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell membrane; Multi-pass membrane protein. Tissue Specificity Cell membrane; Multi-pass membrane protein. Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein-coupled receptor 1 family. Background Offactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arts are responsible for function and goreptin-mediated	Protein Name	Olfactory receptor 52B4 (Olfactory receptor OR11-3)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors share a 7-transmethrane domain structure with many neurotransmitter and hormone receptors sing from single coding-exon genes. Olfactory receptors share a 7-transmethrane domain structure with many neurotransmitter and hormone receptors sind are responsible for the recognition and G protein-mediated	Immunogen	Synthesized peptide derived from human protein . at AA range: 30-110
SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCell membrane; Multi-pass membrane protein.Call PathwayCell membrane; Multi-pass membrane protein.Tissue Specificityfunction:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family.BackgroundOlfactory receptors interact with odorant molecules in the nose, to initiate a neuronal responsible for large family of G-protein coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a -transmembrane domain structure with many neurotransmitter and hormone arising from single coding-exon genes. Olfactory receptors share a -transmembrane domain structure with many neurotransmitter and hormone arising from single coding-exon genes. Olfactory receptors share a -transmembrane domain structure with many neurotransmitter and hormone arising from single coding-exon genes. Olfactory receptors share a -transmembrane domain structure with many neurotransmitter and hormone arising from single coding-exon genes. Olfactory receptors share a -transmembrane domain structure with many neurotransmitter and hormone arising from single coding-exon genes. Olfactory receptor share a -transmetare domain structure with many neuromane transmetared domain structure with many neurotransmitter and	Specificity	O52B4 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americans, similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are reesponsible for the recognition and G protein-mediated	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors arising from single coding-exon genes. Olfactory receptors share a	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from and responsible for the recognition and G protein-mediated	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Cell membrane; Multi-pass membrane protein. Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors g(GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmebrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Concentration	1 mg/ml
Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors arising from single coding excon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Purity	≥90%
Observed Band34kDCell PathwayCell membrane; Multi-pass membrane protein.Tissue SpecificityFunctionfunction:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family.BackgroundOlfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Storago Stability	
Cell PathwayCell membrane; Multi-pass membrane protein.Tissue SpecificityFunctionfunction:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family.BackgroundOlfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Storage Stability	-20°C/1 year
Tissue Specificity Function function:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americanssimilarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated		-20°C/1 year
Functionfunction:Odorant receptorpolymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family.BackgroundOlfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Synonyms	
gene coding for this protein is responsible for functional diversity thus producing pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family.BackgroundOlfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Synonyms Observed Band	34kD
neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Synonyms Observed Band Cell Pathway	34kD
	Synonyms Observed Band Cell Pathway Tissue Specificity	34kD Cell membrane; Multi-pass membrane protein. function:Odorant receptor .,polymorphism:A stop codon at position Ala-40 in the gene coding for this protein is responsible for functional diversity thus producing a

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务, 欢迎咨询



	the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd