



## **OR2L2 Monoclonal Antibody**

2L4)  Immunogen  Synthesized peptide derived from human protein . at AA range: 70-150    Specificity  OR2L2 Monoclonal Antibody detects endogenous levels of protein.    Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.    Source  Monoclonal, Mouse, IgG    Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  -0bserved Band    Observed Band  34kD    Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain,Testis,    Function  offactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Offactory receptors interact with odorant melecules in the nose, to initiate a neuronal response that trigge family of G-protein-coupled receptor (GPCF arising from single coding-exon genes. Olfactory receptors (GPCF arising from single coding-exon genes. Olfactory receptor gene family is the large family for G-protein-coupled receptors (GPCF arising from single coding-exon genes. Olfactory receptors (GPCF arising from single coding-exon genes. Olfactory receptor gene family is the large family for G-protein-coupled receptors (GPCF arising from single coding-exon genes. Olfactory receptors gene family is the large family is the large family for G-protein-coupled receptor (		
ReactivityHuman;Rat;Mouse;ApplicationsWBGene NameOR2L2 OR2L12 OR2L4PProtein NameOlfactory receptor 2L2 (HTPCRH07) (Olfactory receptor 2L12) (Olfactory receptor 2L2)ImmunogenSynthesized peptide derived from human protein . at AA range; 70-150SpecificityOR2L2 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCell membrane; Multi-pass membrane protein.Tissue SpecificityBrain,Testis,Functionfunction:Odorant receptor .,similarity:Belongs to the G-protein coupled recept ramily.,Backgroundolfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptor son ere of a large family of G-protein-coupled receptor arising from single coding-exo genes. Olfactory receptor schere a returnal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exo genes. Olfactory receptor schere a returnal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exo genes. Olfactory receptor schere a returnal membrane	Catalog No	BYmab-07608
Applications  WB    Gene Name  OR2L2 OR2L12 OR2L4P    Protein Name  Olfactory receptor 2L2 (HTPCRH07) (Olfactory receptor 2L12) (Olfactory rece 2L4)    Immunogen  Synthesized peptide derived from human protein . at AA range: 70-150    Specificity  OR2L2 Monoclonal Antibody detects endogenous levels of protein.    Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.    Source  Monoclonal, Mouse, IgG    Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain, Testis,    Function  function:Odorant receptor _,similarity:Belongs to the G-protein coupled recept family.,    Background  Olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smill. The olfactory receptor arising from single coding-exon genes. Olfactory receptor sanar a -7-transmembrane domain structure with many neurotransmilter and hormone receptors and are responsible for the recognition and 6 protein-mediated transducti	Isotype	lgG
Gene Name  OR2L2 OR2L12 OR2L4P    Protein Name  Olfactory receptor 2L2 (HTPCRH07) (Olfactory receptor 2L12) (Olfactory rece 2L4)    Immunogen  Synthesized peptide derived from human protein . at AA range: 70-150    Specificity  OR2L2 Monoclonal Antibody detects endogenous levels of protein.    Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.    Source  Monoclonal, Mouse, IgG    Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain, Testis,    Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,    Background  Olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perceptors of a grote in-coupled receptor arising from single coding-exon genes. Olfactory receptor sanae a -7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responselible for the recognition and grotein-mediated transduction of odorant signals. The olfactory receptor senae a -7-transmembra	Reactivity	Human;Rat;Mouse;
Protein Name  Olfactory receptor 2L2 (HTPCRH07) (Olfactory receptor 2L12) (Olfactory rece 2L4)    Immunogen  Synthesized peptide derived from human protein . at AA range: 70-150    Specificity  OR2L2 Monoclonal Antibody detects endogenous levels of protein.    Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.    Source  Monoclonal, Mouse, IgG    Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  -    Cell Pathway  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain,Testis,    Function  function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor family.,    Background  olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant singles. The olfactory receptor share a 7-transmembrane domain st	Applications	WB
2L4)  Immunogen  Synthesized peptide derived from human protein . at AA range: 70-150    Specificity  OR2L2 Monoclonal Antibody detects endogenous levels of protein.    Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.    Source  Monoclonal, Mouse, IgG    Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  Observed Band    Observed Band  34kD    Cell Pathway  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain,Testis,    Function  offactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant melecules in the nose, to initiate a neuronal response that triggets the perception of a smell. The olifectory receptor growing the a protein or and toporans triggets the perception of a smell. The olifectory receptor growing the parception and G protein-mediated transmerbrane domain structure with many neuroframsmitter and hormone receptors and are responsible for the receptions and G protein-mediated transmerbrane domain structure with many neuroframsmitter and hormone receptors and are responsible for the receptiors and Foropresenter and hormone receptors and are responsible for the reception and G pro	Gene Name	OR2L2 OR2L12 OR2L4P
Specificity  OR2L2 Monoclonal Antibody detects endogenous levels of protein.    Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.    Source  Monoclonal, Mouse, IgG    Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  Observed Band    Observed Band  34kD    Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain, Testis,    Function  offactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCP arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmilter and hormone receptors and are responsible for the recognition and G protein-mediated transmembrane domain structure with many neurotransmilter and hormone receptors and are responsible for the recognition and G protein-mediated transmembrane domain structure with many neurotransmilter and hormone receptors and are responsible for the recognition and G protein-mediated transmembrane domain structure with many neurotransmilter and hormone receptors and are responsible for the recognition and G protein-mediated transmembrane domain	Protein Name	Olfactory receptor 2L2 (HTPCRH07) (Olfactory receptor 2L12) (Olfactory receptor 2L4)
Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.    Source  Monoclonal, Mouse,IgG    Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms	Immunogen	Synthesized peptide derived from human protein . at AA range: 70-150
SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCell membrane; Multi-pass membrane protein.Cill PathwayCell membrane; Multi-pass membrane protein.Tissue SpecificityBrain, Testis,Functionfunction:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,Backgroundolfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptor sant are members of a large family of G-protein-coupled receptor sarien framily.,Backgroundolfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors and are responsible for the receptingion of a smell. The olfactory receptor sarien framily compared of a large family of G-protein-coupled receptor sarien framily compared of a large family of G-protein-coupled receptor sarien framily compared of odorant structure with many neurotransmitter and hormone receptors and are responsible for the receptor genter family to he large transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the receptor family for belarger to of odorant signals. The olfactory receptor genter family is the large transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the receptor genter family is the larger transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the receptor genter family is the larger transmembrane domain structur	Specificity	OR2L2 Monoclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  Observed Band    Observed Band  34kD    Cell Pathway  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain,Testis,    Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptor sinteract with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor sare a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.    Dilution  WB 1:500-2000    Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  -20°C/1 year    Observed Band  34kD    Cell Pathway  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain,Testis,    Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,    Background  Olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens offactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors sand are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Source	Monoclonal, Mouse,IgG
Concentration  1 mg/ml    Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  -20°C/1 year    Observed Band  34kD    Cell Pathway  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain,Testis,    Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,    Background  olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor sinsing from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Purification	
Purity  ≥90%    Storage Stability  -20°C/1 year    Synonyms  -20°C/1 year    Observed Band  34kD    Cell Pathway  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain,Testis,    Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,    Background  olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year    Synonyms  -20°C/1 year    Observed Band  34kD    Cell Pathway  Cell membrane; Multi-pass membrane protein.    Tissue Specificity  Brain, Testis,    Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,    Background  olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Concentration	1 mg/ml
SynonymsObserved Band34kDCell PathwayCell membrane; Multi-pass membrane protein.Tissue SpecificityBrain, Testis,Functionfunction:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,Backgroundolfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Purity	≥90%
Observed Band34kDCell PathwayCell membrane; Multi-pass membrane protein.Tissue SpecificityBrain, Testis,Functionfunction:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,BackgroundOlfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Storage Stability	-20°C/1 year
Cell PathwayCell membrane; Multi-pass membrane protein.Tissue SpecificityBrain, Testis,Functionfunction:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,Backgroundolfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCF) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the larger	Synonyms	
Tissue SpecificityBrain, Testis,Functionfunction:Odorant receptor ., similarity:Belongs to the G-protein coupled recept family.,Backgroundolfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the larger	Observed Band	34kD
Functionfunction:Odorant receptor .,similarity:Belongs to the G-protein coupled recept family.,Backgroundolfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Cell Pathway	Cell membrane; Multi-pass membrane protein.
family., Background olfactory receptor family 2 subfamily L member 2(OR2L2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Tissue Specificity	Brain,Testis,
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory recep proteins are members of a large family of G-protein-coupled receptors (GPCF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large	Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
	Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone

## Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Nanjing BYabscience technology Co.,Ltd