




## OR7A2 Monoclonal Antibody

Catalog No         BYmab-07626           Isotype         IgG           Reactivity         Human;Rat;Mouse;           Applications         WB           Gene Name         OR7A2P OR7A2 OR7A7           Protein Name         Putative olfactory receptor 7A2 (Putative olfactory receptor 7A7)           Immunogen         Synthesized peptide derived from human protein . at AA range: 170-250           Specificity         OR7A2 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         34kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function         caution:Could be the product of a pseudogene., function:Odorant receptor . denominative in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane dom		
Reactivity Human;Rat;Mouse;  Applications WB  Gene Name OR7A2P OR7A2 OR7A7  Protein Name Putative olfactory receptor 7A2 (Putative olfactory receptor 7A7)  Immunogen Synthesized peptide derived from human protein . at AA range: 170-250  Specificity OR7A2 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene, function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family.  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and	Catalog No	BYmab-07626
Applications WB Gene Name OR7A2P OR7A2 OR7A7 Protein Name Putative olfactory receptor 7A2 (Putative olfactory receptor 7A7) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR7A2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:Could be the product of a pseudogene, function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Isotype	IgG
Gene Name         OR7A2P OR7A2 OR7A7           Protein Name         Putative olfactory receptor 7A2 (Putative olfactory receptor 7A7)           Immunogen         Synthesized peptide derived from human protein . at AA range: 170-250           Specificity         OR7A2 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         34kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function         caution:Could be the product of a pseudogene, function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family.           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated	Reactivity	Human;Rat;Mouse;
Protein Name Putative olfactory receptor 7A2 (Putative olfactory receptor 7A7)  Immunogen Synthesized peptide derived from human protein . at AA range: 170-250  Specificity OR7A2 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  Caution:Could be the product of a pseudogene.,function:Odorant receptor  similarity.Belongs to the G-protein coupled receptor 1 family.  Joint and 1 family.  Offactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and the genes and	Applications	WB
Immunogen         Synthesized peptide derived from human protein . at AA range: 170-250           Specificity         OR7A2 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         34kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function         caution:Could be the product of a pseudogene, function:Odorant receptor ,similarity:Belongs to the G-protein coupled receptor 1 family.,           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and the genes and	Gene Name	OR7A2P OR7A2 OR7A7
Specificity OR7A2 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene, function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and	Protein Name	Putative olfactory receptor 7A2 (Putative olfactory receptor 7A7)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene.,function:Odorant receptorsimilarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and	Immunogen	Synthesized peptide derived from human protein . at AA range: 170-250
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         34kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function         caution:Could be the product of a pseudogene, function:Odorant receptor, similarity:Belongs to the G-protein coupled receptor 1 family.           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triaggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Specificity	OR7A2 Monoclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  34kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Purification	•
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene.,function:Odorant receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Dilution	WB 1:500-2000
Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Concentration	1 mg/ml
Synonyms  Observed Band 34kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Purity	≥90%
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Storage Stability	-20°C/1 year
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Synonyms	
Tissue Specificity  Function  caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Observed Band	34kD
Function  caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Tissue Specificity	
neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and	Function	

Nanjing BYabscience technology Co.,Ltd







RefSeq, Jul 2008],

matters needing	
attention	

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

