



TRXR1 Monoclonal Antibody

Catalog No	BYmab-07663
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TXNRD1 GRIM12 KDRF
Protein Name	Thioredoxin reductase 1, cytoplasmic (TR) (EC 1.8.1.9) (Gene associated with retinoic and interferon-induced mortality 12 protein) (GRIM-12) (Gene associated with retinoic and IFN-induced mortality 12
Immunogen	Synthesized peptide derived from part region of human protein AA range: 268-318
Specificity	TRXR1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	71kD
Cell Pathway	[Isoform 1]: Cytoplasm .; [Isoform 4]: Cytoplasm . Nucleus .; [Isoform 5]: Cytoplasm .
Tissue Specificity	[Isoform 1]: Expressed predominantly in Leydig cells (at protein level). Also expressed in ovary, spleen, heart, liver, kidney and pancreas and in a number of cancer cell lines. ; [Isoform 4]: Widely expressed with highest levels in kidney, testis, uterus, ovary, prostate, placenta and fetal liver.
Function	catalytic activity:Thioredoxin + NADP(+) = thioredoxin disulfide + NADPH.,cofactor:Binds 1 FAD per subunit.,domain:The N-terminal glutaredoxin domain found in isoform 1 does not contain the C-P-Y-C redox-active motif normally found in glutaredoxins and has been found to be inactive in classical glutaredoxin assays.,function:Isoform 1 may possess glutaredoxin activity as well as thioredoxin reductase activity and induces actin and tubulin polymerization, leading to formation of cell membrane protrusions. Isoform 4 enhances the

Nanjing BYabscience technology Co.,Ltd



transcriptional activity of estrogen receptors alpha and beta while isoform 5 enhances the transcriptional activity of the beta receptor only. Isoform 5 also mediates cell death induced by a combination of interferon-beta and retinoic acid.,induction:Isoform 5 is induced by a combination of interferon-beta and retinoic acid (at protein level). Isoform 1 is induced b

Background

thioredoxin reductase 1(TXNRD1) Homo sapiens This gene encodes a member of the family of pyridine nucleotide oxidoreductases. This protein reduces thioredoxins as well as other substrates, and plays a role in selenium metabolism and protection against oxidative stress. The functional enzyme is thought to be a homodimer which uses FAD as a cofactor. Each subunit contains a selenocysteine (Sec) residue which is required for catalytic activity. The selenocysteine is encoded by the UGA codon that normally signals translation termination. The 3' UTR of selenocysteine-containing genes have a common stem-loop structure, the sec insertion sequence (SECIS), that is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. Alternative splicing results in several transcript variants encoding the same or different isoforms. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658