



COPB Monoclonal Antibody

Catalog No	BYmab-07670	
Isotype	IgG	
Reactivity	Human;Rat;Mouse;	
Applications	WB	
Gene Name	COPB1 COPB MSTP026	
Protein Name	Coatomer subunit beta (Beta-coat protein) (Beta-COP)	
Immunogen	Synthesized peptide derived from part region of human protein AA range: 865-915	
Specificity	COPB Monoclonal Antibody detects endogenous levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Source	Monoclonal, Mouse,IgG	
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution	WB 1:500-2000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Observed Band	104kD	
Cell Pathway	Cytoplasm. Golgi apparatus membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, COPI-coated vesicle membrane; Peripheral membrane protein ; Cytoplasmic side . Cell membrane . Endoplasmic reticulum-Golgi intermediate compartment . The coatomer is cytoplasmic or polymerized on the cytoplasmic side of the Golgi, as well as on the vesicles/buds originating from it (By similarity). Proteolytic cleavage by CAPN8 triggers translocation from Golgi to cytoplasm (By similarity). Found in perinuclear vesicular-tubular clusters (VTCs) and in the Golgi region where associated with vesicles, buds and rims of the Golgi stack (By similarity). Occasionally present at the trans-side of Golgi, but mainly present at the cis-Golgi side in transitional areas (TA), on so-called periph	
Tissue Specificity	Amygdala,Aorta,Lymphoid,Testis,	
Function	function:The coatomer is a cytosolic protein complex that binds to dilysine motifs and reversibly associates with Golgi non-clathrin-coated vesicles, which further	
Nanjing BYabscience technology Co.,Ltd		

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658

K	博研生物
	BYabscience

国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务,欢迎咨询



	mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins; the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors.,miscellaneous:Brefeldin A induces dissociation from the Golgi of the beta-COP and presumably the other coatomer subunits.,PTM:Proteolytically cleaved between Ser-528 and Ser-529 by CAPN8.,subcellular locat	
Background	This gene encodes a protein subunit of the coatomer complex associated with non-clathrin coated vesicles. The coatomer complex, also known as the coat protein complex 1, forms in the cytoplasm and is recruited to the Golgi by activated guanosine triphosphatases. Once at the Golgi membrane, the coatomer complex may assist in the movement of protein and lipid components back to the endoplasmic reticulum. Alternatively spliced transcript variants have been described. [provided by RefSeq, Jan 2009],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
Products Images		

Nanjing BYabscience technology Co.,Ltd