




## DC1L1 Monoclonal Antibody

which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC		
Reactivity         Human;Rat;Mouse;           Applications         WB           Gene Name         DYNC1LI1 DNCLI1           Protein Name         Cytoplasmic dynein 1 light intermediate chain 1 (LIC1) (Dynein light chain A) (DLC-A) (Dynein light intermediate chain 1, cytosolic)           Immunogen         Synthesized peptide derived from part region of human protein AA range: 292-342           Specificity         DC1L1 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         57kD           Cell Pathway         Cytoplasm. Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole.           Tissue Specificity         Cerebellum, Coronary artery, Epithelium, Human testis,           Function         function: May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity. PTM: Phosphorylated, similarity, selongs	Catalog No	BYmab-07682
Applications  Gene Name  DYNC1LI1 DNCLI1  Protein Name  Cytoplasmic dynein 1 light intermediate chain 1 (LIC1) (Dynein light chain A) (DLC-A) (Dynein light intermediate chain 1, cytosolic)  Immunogen  Synthesized peptide derived from part region of human protein AA range: 292-342  Specificity  DC1L1 Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  57kD  Cell Pathway  Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity  Cerebellum, Coronary artery, Epithelium, Human testis,  function: May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity, PTM-Phosphorylated .smilarity selongs to the dynein light intermediate chain family, subunit-Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  The protein encoded by this gene belongs to light intermediate austicult family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Isotype	IgG
Gene Name DYNC1LI1 DNCLI1  Protein Name Cytoplasmic dynein 1 light intermediate chain 1 (LIC1) (Dynein light chain A) (DLC-A) (Dynein light intermediate chain 1, cytosolic)  Immunogen Synthesized peptide derived from part region of human protein AA range: 292-342  Specificity DC1L1 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms  Observed Band 57kD  Cell Pathway Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity Cerebellum, Coronary artery, Epithelium, Human testis,  Function function: May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity, PTM: Phosphorylated, similarity: Belongs to the dynein light intermediate chain family, subunit: Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Reactivity	Human;Rat;Mouse;
Protein Name Cytoplasmic dynein 1 light intermediate chain 1 (LIC1) (Dynein light chain A) (DLC-A) (Dynein light intermediate chain 1, cytosolic) Immunogen Synthesized peptide derived from part region of human protein AA range: 292-342 Specificity DC1L1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole . Tissue Specificity Cerebellum,Coronary artery,Epithelium,Human testis, Function function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity, PTM:Phosphorylated,.similarity:Belongs to the dynein light intermediate chain family, subunit:Consist of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains., The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome sergegation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC)	Applications	WB
Immunogen         Synthesized peptide derived from part region of human protein AA range: 292-342           Specificity         DC1L1 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         57kD           Cell Pathway         Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .           Tissue Specificity         Cerebellum,Coronary artery,Epithelium,Human testis,           Function         function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM.Phosphorylated, similarity.Belongs to the dynein light intermediate chain family, subunit;Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,           Background         The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and ch	Gene Name	DYNC1LI1 DNCLI1
Immunogen         Synthesized peptide derived from part region of human protein AA range: 292-342           Specificity         DC1L1 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         57kD           Cell Pathway         Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .           Tissue Specificity         Cerebellum,Coronary artery,Epithelium,Human testis,           Function         function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM.Phosphorylated, similarity.Belongs to the dynein light intermediate chain family, subunit;Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,           Background         The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and ch	Protein Name	Cytoplasmic dynein 1 light intermediate chain 1 (LIC1) (Dynein light chain A) (DLC-A) (Dynein light intermediate chain 1, cytosolic)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 57kD  Cell Pathway Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity Cerebellum, Coronary artery, Epithelium, Human testis,  Function function: May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity. PTM: Phosphorylated., similarity. Belongs to the dynein light intermediate chain family., subunit: Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Immunogen	Synthesized peptide derived from part region of human protein AA range:
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band       57kD         Cell Pathway       Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .         Tissue Specificity       Cerebellum, Coronary artery, Epithelium, Human testis,         Function       function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity, PTM:Phosphorylated, similarity:Belongs to the dynein light intermediate chain family, subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.         Background       The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Specificity	DC1L1 Monoclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  57kD  Cell Pathway  Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity  Cerebellum,Coronary artery,Epithelium,Human testis,  Function  function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family.,subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 57kD  Cell Pathway Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity Cerebellum,Coronary artery,Epithelium,Human testis,  Function function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated, similarity:Belongs to the dynein light intermediate chain family, subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.  Background The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 57kD  Cell Pathway Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity Cerebellum,Coronary artery,Epithelium,Human testis,  Function function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family.,subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 57kD  Cell Pathway Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity Cerebellum,Coronary artery,Epithelium,Human testis,  Function function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family.,subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year  Synonyms  Observed Band  57kD  Cell Pathway  Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity  Cerebellum, Coronary artery, Epithelium, Human testis,  function: May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity., PTM:Phosphorylated., similarity:Belongs to the dynein light intermediate chain family., subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Concentration	1 mg/ml
Synonyms  Observed Band  57kD  Cell Pathway  Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity  Cerebellum,Coronary artery,Epithelium,Human testis,  function  function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family, subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC)	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Tissue Specificity  Cerebellum, Coronary artery, Epithelium, Human testis,  function: May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity., PTM: Phosphorylated., similarity: Belongs to the dynein light intermediate chain family., subunit: Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC)	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole .  Cerebellum, Coronary artery, Epithelium, Human testis,  function  function: May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity., PTM: Phosphorylated., similarity: Belongs to the dynein light intermediate chain family., subunit: Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC)	Synonyms	
Tissue Specificity  Cerebellum,Coronary artery,Epithelium,Human testis,  function  function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family.,subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC)	Observed Band	57kD
Function  function:May play a role in binding dynein to membranous organelles or chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family.,subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC)	Cell Pathway	
chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family.,subunit:Consists of at least 3 heavy chains (alpha, beta and gamma), 2 intermediate chains and 8 light chains.,  Background  The protein encoded by this gene belongs to light intermediate subunit family, whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC)	Tissue Specificity	Cerebellum, Coronary artery, Epithelium, Human testis,
whose members are components of the multiprotein cytoplasmic dynein complex which is involved in intracellular trafficking and chromosome segregation during mitosis. The protein plays a role in moving the spindle assembly checkpoint (SAC	Function	chromosomes. May associate with the heavy chains in the dynein head where they might regulate enzyme activity.,PTM:Phosphorylated.,similarity:Belongs to the dynein light intermediate chain family.,subunit:Consists of at least 3 heavy
	Background	whose members are components of the multiprotein cytoplasmic dynein complex,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	cargo molecules to facilitate intracellular vesicle trafficking. [provided by RefSeq, Jul 2016],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd