



RBP10 Monoclonal Antibody

Catalog No	BYmab-07686
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	RANBP10 KIAA1464
Protein Name	Ran-binding protein 10 (RanBP10)
Immunogen	Synthesized peptide derived from part region of human protein AA range: 353-403
Specificity	RBP10 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
	0010/4
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year 68kD
Synonyms	
Synonyms Observed Band	68kD
Synonyms Observed Band Cell Pathway	68kD Cytoplasm, cytosol . Nucleus . Predominantly cytoplasmic.
Synonyms Observed Band Cell Pathway Tissue Specificity	68kD Cytoplasm, cytosol . Nucleus . Predominantly cytoplasmic. Broadly expressed, with highest levels in skeletal muscle. domain:The SPRY domain mediates the interaction with MET.,function:May act as an adapter protein to couple membrane receptors to intracellular signaling pathways. In contrast to RANBP9, does not interact with Sos and does not activate the Ras pathway.,similarity:Belongs to the RANBP9/10 family.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 CTLH domain.,similarity:Contains 1 LisH domain.,subunit:Interacts with RAN. Interacts with the HGF receptor MET.,tissue specificity:Broadly expressed, with highest

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务,欢迎咨询



Products Images	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
matters needing attention	Avoid repeated freezing and thawing!
	encoded by this gene is a cytoplasmic guanine nucleotide exchange factor (GEF) that binds beta-tubulin and has GEF activity toward RAN. The encoded protein plays a role in the formation of noncentrosomal microtubules. In addition, this protein may be involved in the regulation of D(1) receptor signaling by protein kinase C delta and protein kinase C gamma. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2016],

Nanjing BYabscience technology Co.,Ltd