



LIMA1 Monoclonal Antibody

Catalog No	BYmab-07687
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	LIMA1 EPLIN SREBP3 PP624
Protein Name	LIM domain and actin-binding protein 1 (Epithelial protein lost in neoplasm)
Immunogen	Synthesized peptide derived from part region of human protein AA range: 631-681
Specificity	LIMA1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	83kD
Cell Pathway	Cytoplasm. Cell junction, focal adhesion . Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, stress fiber . Cell membrane . Expressed in the brush border membrane of the small intestine and colocalizes with NPC1L1 and MYO5B (PubMed:29880681). Colocalizes with PXN at focal adhesions in mesangial cells (PubMed:24694988). Colocalizes with actin stress fibers in quiescent cells. PDGF stimulation induced disassembly of stress fibers and formation of peripheral and dorsal ruffles, where LIMA1 is relocalized (By similarity). .
Tissue Specificity	Highly expressed in placenta, kidney, pancreas, prostate, ovary, spleen and heart. Also detected in lung, liver, brain, skeletal muscle, thymus, testis and intestine. Not detected in leukocytes. Isoform Beta expressed generally at very low levels. Isoform Alpha abundant in epithelial cells from mammary gland, prostate and in normal oral keratinocytes. Low levels in aortic endothelial cells and dermal fibroblasts. Not detectable in myocardium.
Function	domain:Contains at least 2 actin-binding domains, one on each side of the LIM domain. Both domains bind actin monomers and filaments. The C-terminal

Nanjing BYabscience technology Co.,Ltd



domain binds filaments more efficiently than the N-terminus.,function: Binds to actin monomers and filaments. Increases the number and size of actin stress fibers and inhibits membrane ruffling. Inhibits actin filament depolymerization. Bundles actin filaments, delays filament nucleation and reduces formation of branched filaments.,induction: Down-regulated in some cancer cell lines. Isoform Alpha induced by serum. Isoform Beta constitutively expressed.,similarity: Contains 1 LIM zinc-binding domain.,subcellular location: This cytoskeletal protein co-localizes with actin stress fibers and focal adhesion plaques.,tissue specificity: Highly expressed in placenta, kidney, pancreas, prostate, ovary, spleen and heart. Also detected in lung, liver, bra

Background

This gene encodes a cytoskeleton-associated protein that inhibits actin filament depolymerization and cross-links filaments in bundles. It is downregulated in some cancer cell lines. Alternatively spliced transcript variants encoding different isoforms have been described for this gene, and expression of some of the variants maybe independently regulated. [provided by RefSeq, Aug 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images