



TIAM1 Monoclonal Antibody

Catalog No	BYmab-07722
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TIAM1
Protein Name	T-lymphoma invasion and metastasis-inducing protein 1 (TIAM-1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TIAM1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	175kD
Cell Pathway	Cell junction. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Detected at the boundary between cells with actin-rich protrusions (By similarity). Presence of KRIT1, CDH5 and RAP1B is required for its localization to the cell junction. .
Tissue Specificity	Found in virtually all analyzed tumor cell lines including B- and T-lymphomas, neuroblastomas, melanomas and carcinomas.
Function	function:Modulates the activity of RHO-like proteins and connects extracellular signals to cytoskeletal activities. Acts as a GDP-dissociation stimulator protein that stimulates the GDP-GTP exchange activity of RHO-like GTPases and activates them. Activates RAC1, CDC42, and to a lesser extent RHOA.,similarity:Belongs to the TIAM family.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 RBD (Ras-binding) domain.,similarity:Contains 2 PH domains.,subunit:Interacts with BAIAP2.,tissue specificity:Found in virtually all analyzed tumor cell lines including B- and

Nanjing BYabscience technology Co.,Ltd



T-lymphomas, neuroblastomas, melanomas and carcinomas.,

Background

function:Modulates the activity of RHO-like proteins and connects extracellular signals to cytoskeletal activities. Acts as a GDP-dissociation stimulator protein that stimulates the GDP-GTP exchange activity of RHO-like GTPases and activates them. Activates RAC1, CDC42, and to a lesser extent RHOA.,similarity:Belongs to the TIAM family.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 RBD (Ras-binding) domain.,similarity:Contains 2 PH domains.,subunit:Interacts with BAIAP2.,tissue specificity:Found in virtually all analyzed tumor cell lines including B- and T-lymphomas, neuroblastomas, melanomas and carcinomas.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658