



COAA1 Monoclonal Antibody

Catalog No	BYmab-07735
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	COL10A1
Protein Name	Collagen alpha-1(X) chain
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	COAA1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	74kD
Cell Pathway	Secreted, extracellular space, extracellular matrix .
Tissue Specificity	
Function	disease:Defects in COL10A1 are the cause of Schmid type metaphyseal chondrodysplasia (SMCD) [MIM:156500]. SMCD is a dominantly inherited

chondrodysplasia (SMCD) [MIM:156500]. SMCD is a dominantly inherited disorder of the osseous skeleton. The cardinal features of the phenotype are mild short stature, coxa vara and a waddling gait. Radiography usually shows sclerosis of the ribs, flaring of the metaphyses, and a wide irregular growth plate, especially of the knees. A variant form of SMCD is spondylometaphyseal dysplasia Japanese type. It is characterized by spinal involvement comprising mild platyspondyly, vertebral body abnormalities, and end-plate irregularity.,function:Type X collagen is a product of hyperthrophic chondrotocytes and has been localized to presumptive mineralization zones of hyaline cartilage.,PTM:Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.,similarity:Contains 1 C

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	This gene encodes the alpha chain of type X collagen, a short chain collagen expressed by hypertrophic chondrocytes during endochondral ossification. Unlike type VIII collagen, the other short chain collagen, type X collagen is a homotrimer. Mutations in this gene are associated with Schmid type metaphyseal chondrodysplasia (SMCD) and Japanese type spondylometaphyseal dysplasia (SMD). [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd