



NUCG Monoclonal Antibody

Catalog No	BYmab-07798
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	ENDOG
Protein Name	Endonuclease G, mitochondrial (Endo G) (EC 3.1.30)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	NUCG Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,lgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration	i nig/ini
Purity	≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 32kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 32kD Mitochondrion . Brain,Uterus, cofactor:Manganese or magnesium.,function:Cleaves DNA at double-stranded (DG)n.(DC)n and at single-stranded (DC)n tracts. In addition to deoxyribonuclease activities, also has ribonuclease (RNase) and RNase H activities. CaMABle of generating the RNA primers required by DNA polymerase gamma to initiate replication of mitochondrial DNA.,miscellaneous:The active site
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 32kD Mitochondrion . Brain,Uterus, cofactor:Manganese or magnesium.,function:Cleaves DNA at double-stranded (DG)n.(DC)n and at single-stranded (DC)n tracts. In addition to deoxyribonuclease activities, also has ribonuclease (RNase) and RNase H activities. CaMABle of generating the RNA primers required by DNA polymerase gamma to initiate replication of mitochondrial DNA.,miscellaneous:The active site contains 1 hydrated divalent metal cation that has only 1 direct interaction with the protein; all other interactions are via water molecules.,similarity:Belongs to the

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	RNA primers required by DNA polymerase gamma to initiate replication of mitochondrial DNA. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658