



HIC2 Monoclonal Antibody

Catalog No	BYmab-07808
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	HIC2 HRG22 KIAA1020 ZBTB30
Protein Name	Hypermethylated in cancer 2 protein (Hic-2) (HIC1-related gene on chromosome 22 protein) (Hic-3) (Zinc finger and BTB domain-containing protein 30)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HIC2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	67kD
Cell Pathway	Nucleus.
Tissue Specificity	Highest levels in cerebellum.
Function	function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family. Hic subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Interacts with CtBP.,tissue specificity:Highest levels in cerebellum.,
Background	function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family. Hic subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Interacts with CtBP.,tissue specificity:Highest levels in cerebellum.,
matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images