



ILF2 Monoclonal Antibody

Catalog No	BYmab-07814
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ILF2 NF45 PRO3063
Protein Name	Interleukin enhancer-binding factor 2 (Nuclear factor of activated T-cells 45 kDa)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ILF2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	42kD
Cell Pathway	Nucleus, nucleolus. Cytoplasm . Nucleus . Localized in cytoplasmic mRNP granules containing untranslated mRNAs.
Tissue Specificity	Fetal liver,Lung,Thymus,
Function	function:Appears to function predominantly as a heterodimeric complex with ILF3. This complex may regulate transcription of the IL2 gene during T-cell activation. It can also promote the formation of stable DNA-dependent protein kinase holoenzyme complexes on DNA.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Chimeric cDNA.,similarity:Contains 1 DZF domain.,subunit:Forms heterodimers with ILF3. ILF2-ILF3 heterodimers may also bind to PRKDC/XRCC7: this may stabilize the interaction of PRKDC/XRCC7 and the heterodimeric complex of G22P1/KU70 and XRCC5/KU80.,
Background	The protein encoded by this gene is a transcription factor required for T-cell expression of the interleukin 2 gene. It also binds RNA and is an essential component for encapsidation and protein priming of hepatitis B viral polymerase.

Nanjing BYabscience technology Co.,Ltd



The encoded 45 kDa protein (NF45, ILF2) forms a complex with the 90 kDa interleukin enhancer-binding factor 3 (NF90, ILF3), and this complex has been shown to affect the redistribution of nuclear mRNA to the cytoplasm, to repair DNA breaks by nonhomologous end joining, and to negatively regulate the microRNA processing pathway. Knockdown of NF45 or NF90 protein retards cell growth, possibly by inhibition of mRNA stabilization. Alternative splicing results in multiple transcript variants. Related pseudogenes have been found on chromosomes 3 and 14. [provided by RefSeq, Dec 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ILF2 Monoclonal Antibody