



VAMP7 Monoclonal Antibody

Catalog No	BYmab-07842
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	VAMP7 SYBL1
Protein Name	Vesicle-associated membrane protein 7 (VAMP-7) (Synaptobrevin-like protein 1) (Tetanus-insensitive VAMP) (Ti-VAMP)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	VAMP7 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	24kD
Cell Pathway	Cytoplasmic vesicle, secretory vesicle membrane ; Single-pass type IV membrane protein . Golgi apparatus, trans-Golgi network membrane ; Single-pass type IV membrane protein . Late endosome membrane ; Single-pass type IV membrane protein . Lysosome membrane ; Single-pass type IV membrane protein . Endoplasmic reticulum membrane ; Single-pass type IV membrane protein . Cytoplasmic vesicle, phagosome membrane ; Single-pass type IV membrane protein . Cell junction, synapse, synaptosome . In immature neurons expression is localized in vesicular structures in axons and dendrites while in mature neurons it is localized to the somatodendritic region. Colocalizes with LAMP1 in kidney cells. Localization to the endoplasmic reticulum membrane was observed in the intestine but not in liver or kidney
Tissue Specificity	Detected in all tissues tested.
Function	function:Involved in the targeting and/or fusion of transport vesicles to their target membrane during transport of proteins from the early endosome to the lysosome.

Nanjing BYabscience technology Co.,Ltd



Required for heterotypic fusion of late endosomes with lysosomes and homotypic lysosomal fusion. Required for calcium regulated lysosomal exocytosis. Involved in the export of chylomicrons from the endoplasmic reticulum to the cis Golgi. Required for exocytosis of mediators during eosinophil and neutrophil degranulation, and target cell killing by natural killer cells. Required for focal exocytosis of late endocytic vesicles during phagosome formation.,miscellaneous:Loss-of-function mutant (antisense inhibition) displays impaired granzyme B release and target cell killing by natural killer cells.,miscellaneous:The gene encoding for this protein is located in the pseudoautosomal region 2 (PAR2) of X and Y chromosomes.,similar

Background

This gene encodes a transmembrane protein that is a member of the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) family. The encoded protein localizes to late endosomes and lysosomes and is involved in the fusion of transport vesicles to their target membranes. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Jun 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using VAMP7 Monoclonal Antibody