



DERL2 Monoclonal Antibody


Catalog No	BYmab-07866
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DERL2 DER2 FLANA CGI-101 SBBI53
Protein Name	Derlin-2 (Degradation in endoplasmic reticulum protein 2) (DERtrin-2) (Der1-like protein 2) (F-LAN-1) (F-LANa)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DERL2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	26kD
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
Tissue Specificity	Ubiquitous. Overexpressed in various hepatocarcinomas.
Function	function:Functional component of endoplasmic reticulum-associated degradation (ERAD) for misfolded luminal glycoproteins, but not that of misfolded nonglycoproteins. May act by forming a channel that allows the retrotranslocation of misfolded glycoproteins into the cytosol where they are ubiquitinated and degraded by the proteasome. May mediate the interaction between VCP and the degradation substrate. In contrast to DERL1, it is not involved in the degradation of MHC class I heavy chains following infection by cytomegaloviruses. May play a role in cell proliferation.,induction:Up-regulated in response to ER stress via the ERN1-XBP1 pathway of the unfolded protein response (UPR).,similarity:Belongs to the derlin family.,subunit:Forms homo- and heterooligomers with DERL3 and, to a lesser extent, with DERL1. Interacts with SELS, VCP and EDEM1. Mediates association between VCP and EDEM1, as

Nanjing BYabscience technology Co.,Ltd



Background	derlin 2(DERL2) Homo sapiens Proteins that are unfolded or misfolded in the endoplasmic reticulum (ER) must be refolded or degraded to maintain the homeostasis of the ER. DERL2 is involved in the degradation of misfolded glycoproteins in the ER (Oda et al., 2006 [PubMed 16449189]).[supplied by OMIM, Mar 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using DERL2 Monoclonal Antibody