



KLD8B mouse mAb

Catalog No BYmab-07942 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name KLHDC8B FP17659 Protein Name KLD8B Immunogen Synthesized peptide derived from human KLD8B AA range: 120-170 Specificity This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse,lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm: Midbody. In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytoplasmic bridge linking daughter cells as they are about to separate during cytoplasmic bridge linking daughter cells as they are about to separate during cytoplasmic bridge linking daughter cells as they are about to separate during cytoplasmic bridge linking daug		
Reactivity Human; Mouse;Rat Applications WB Gene Name KLHDC8B FP17659 Protein Name KLD8B Immunogen Synthesized peptide derived from human KLD8B AA range: 120-170 Specificity This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm. Midbody. In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex, Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Catalog No	BYmab-07942
Applications WB Gene Name KLHDC8B FP17659 Protein Name KLD8B Immunogen Synthesized peptide derived from human KLD8B AA range: 120-170 Specificity This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex, Brain, Function similarity: Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name KLHDC8B FP17659 Protein Name KLD8B Immunogen Synthesized peptide derived from human KLD8B AA range: 120-170 Specificity This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Reactivity	Human; Mouse;Rat
Protein Name KLD8B Immunogen Synthesized peptide derived from human KLD8B AA range: 120-170 Specificity This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis . Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Applications	WB
Immunogen Synthesized peptide derived from human KLD8B AA range: 120-170 Specificity This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm. Midbody. In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Gene Name	KLHDC8B FP17659
Specificity This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm. Midbody. In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Protein Name	KLD8B
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm. Midbody. In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex,Brain, Function Similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human KLD8B AA range: 120-170
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis . Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of KLD8B at Human/Mouse/Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm : Midbody : In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis . Tissue Specificity Adrenal cortex,Brain, Function Similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010]. matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.56% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Kelch domain-containing protein 8B Observed Band 38kD Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis. Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Midbody . In mitotic cells, concentrates in the midbody of the cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis Tissue Specificity Adrenal cortex,Brain, Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Synonyms	Kelch domain-containing protein 8B
cytoplasmic bridge linking daughter cells as they are about to separate during cytokinesis Tissue Specificity Adrenal cortex,Brain, Function Similarity:Contains 8 Kelch repeats., This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Observed Band	38kD
Function similarity:Contains 8 Kelch repeats., Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Cell Pathway	cytoplasmic bridge linking daughter cells as they are about to separate during
Background This gene encodes a protein which forms a distinct beta-propeller protein structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], Matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Adrenal cortex,Brain,
structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq, Sep 2010], matters needing Avoid repeated freezing and thawing!	Function	similarity:Contains 8 Kelch repeats.,
	Background	structure of kelch domains allowing for protein-protein interactions. Mutations in this gene have been associated with Hodgkin lymphoma. [provided by RefSeq,
		Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using KLD8B mouse mAb

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658