



SPTC2 mouse mAb

Catalog NoBYmab-07986IsotypeIgGReactivityHuman; MouseApplicationsWBGene NameSPTLC2 KIAA0526 LCB2Protein NameSPTC2ImmunogenSynthesized peptide derived from human SPTC2 At range: 273-323SpecificityThis antibody detects endogenous levels of SPTC2 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide.SourceMonocional, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionVB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSerine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis protei 2) (LCB 2) (Long chain base biosynthesis protei 2) (LCB2) (LCB 2) (Long chain base biosynthesis protei 2) (LCB2) (LCB 2) (LOG thain base biosynthesis protei 2) (LCB2) (LCB 2) (Long chain base biosynthesis protei 2) (LCB2) (LCB 2) (LOG thain base biosynthesis protei 2) (LCB2) (LCB 2) (LCD chain base biosynthesis protei 2) (LCB2) (LCB 2) (LCD chain base biosynthesis protei 2) (LCB2) (LCB 2) (LOG thain base biosynthesis protei 2) (LCB2) (LCB 2) (LOG chain base biosynthesis protei 2) (LCB2) (LCB 2) (LCD chain base biosynthesis protei 2) (LCB2) (LCB 2) (LCD chain base biosynthesis protei 2) (LCB2) (LCB 2) (LCD chain base biosynthesis protei 2) (LCB2) (LCB 2) (LOG chain base biosynthesis protei 2) (LCB2) (LCB 2) (LCD chain base biosynthesis protei 2) (LCB2) (LCB 2) (LCD chain base biosynthesis protei 2) (LCB2) (LCB 2) (LOB chain base biosynthesis protei 2) (LCB2) (LCB 2)		
ReactivityHuman; MouseApplicationsWBGene NameSPTLC2 KIAA0526 LCB2Protein NameSPTC2ImmunogenSynthesized peptide derived from human SPTC2 AA range: 273-323SpecificityThis antibody detects endogenous levels of SPTC2 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSerine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + co/(2), cofactor:Pyridoxal phosphate, pathway/Lipid metabolism; sphinglipid aminotransferase family, subunit.SPTLC2 and SPTLC3 may encode subunits of the enzyme, tissue specificity/Widely expressed.BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphinglipid biosynthesis. It catalyzes the	Catalog No	BYmab-07986
Applications WB Gene Name SPTLC2 KIAA0526 LCB2 Protein Name SPTC2 Immunogen Synthesized peptide derived from human SPTC2 AA range: 273-323 Specificity This antibody detects endogenous levels of SPTC2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2) Observed Band 60kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass membrane protein . Tissue Specificity Widely expressed. Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor/Pyridoxal phosphatepathway.Lipid metabolism, sphingalipid biosynthesis. It catalyzes the Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphingalipid metabolism, sphingalipid biosynthesis. It catalyzes the </th <td>Isotype</td> <td>IgG</td>	Isotype	IgG
Gene NameSPTLC2 KIAA0526 LCB2Protein NameSPTC2ImmunogenSynthesized peptide derived from human SPTC2 AA range: 273-323SpecificityThis antibody detects endogenous levels of SPTC2 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSerine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane : Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphate, apathway: Lipid metabolism, similarity: Belongs to the class-11 pyridoxal phosphate-dependent aminotransferase family. subunit: SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme, tissue specificity:Widely expressed.BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid	Reactivity	Human; Mouse
Protein NameSPTC2ImmunogenSynthesized peptide derived from human SPTC2 AA range: 273-323SpecificityThis antibody detects endogenous levels of SPTC2 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSerine palmitoytransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LOB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphate, pathway:Lipid metabolism, similarity: Belongs to the class-11 pyridoxal phosphate-dependent aminotransferase familysubunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme, tissue specificity:Widely expressed.BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid	Applications	WB
ImmunogenSynthesized peptide derived from human SPTC2 AA range: 273-323SpecificityThis antibody detects endogenous levels of SPTC2 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSerine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphate, pathway:Lipid metabolism, sphingolipid metabolism, similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase familysuburit.SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme, tissue specificity.Widely expressed.BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid play suburits. It catalyzes the	Gene Name	SPTLC2 KIAA0526 LCB2
Specificity This antibody detects endogenous levels of SPTC2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2) Observed Band 60kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass membrane protein . Tissue Specificity Widely expressed. Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2).cofactor:Pyridoxal phosphate.pathway:Lipd metabolism: sphingolipid store as subunits of the enzyme, tissue specificity:Widely expressed. Background This gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase. Serine palmitoyltransferase. S	Protein Name	SPTC2
FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSerine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO2), cofactor:Pyridoxal phosphate, pathway:Lipid metabolism, sphingolipid metabolism, similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.subunit:SPTLC2 and SPTLC2 and SPTLC2 anay encode subunits of the enzyme, tissue specificity:Widely expressed.BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine p	Immunogen	Synthesized peptide derived from human SPTC2 AA range: 273-323
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis protein 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2) Observed Band 60kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass membrane protein . Tissue Specificity Widely expressed. Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphatepathway:Lipid metabolism; sphingolipid metabolism; similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase familysubunit:SPTLC1, and SPTLC2 and SPTLC3 may encode subunits of the enzymetissue specificity:Widely expressed., Background This gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, to wind the consist of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Specificity	This antibody detects endogenous levels of SPTC2 at Human/Mouse
PurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSerine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2),.cofactor:Pyridoxal phosphatepathway:Lipid metabolism; sphingolipid metabolism,.similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.subunit:SPTLC1 and SPTLC2 and SPTLC3 may encode subunits of the enzyme.tissue specificity:Widely expressed.BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consist of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.100% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2) Observed Band 60kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass membrane protein . Tissue Specificity Widely expressed. Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism, similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family, subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed., Background This gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis protein 2) (LCB 2) (LOB g chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2) Observed Band 60kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass membrane protein . Tissue Specificity Widely expressed. Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphate.apathway:Lipid metabolism, similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase familysubunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed., Background This gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis protein 2) (LCB 2) (LOng chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2) Observed Band 60kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass membrane protein . Tissue Specificity Widely expressed. Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2)., cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism, similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family., subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme., tissue specificity:Widely expressed. Background This gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Serine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis protei 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2) Observed Band 60kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass membrane protein . Tissue Specificity Widely expressed. Function catalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism, similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family, subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme, tissue specificity:Widely expressed. Background This gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Concentration	1 mg/ml
SynonymsSerine palmitoyltransferase 2 (EC 2.3.1.50) (Long chain base biosynthesis prote 2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2), cofactor:Pyridoxal phosphate.pathway:Lipid metabolism; sphingolipid metabolism.,similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.,subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed.,BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Purity	≥90%
2) (LCB 2) (Long chain base biosynthesis protein 2a) (LCB2a) (Serine-palmitoyl-CoA transferase 2) (SPT 2)Observed Band60kDCell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2).,cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism.,similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.,subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed.,BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Storage Stability	-20°C/1 year
Cell PathwayEndoplasmic reticulum membrane ; Single-pass membrane protein .Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2).,cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism.,similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.,subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed.,BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Synonyms	
Tissue SpecificityWidely expressed.Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2).,cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism.,similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.,subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed.,BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Observed Band	60kD
Functioncatalytic activity:Palmitoyl-CoA + L-serine = CoA + 3-dehydro-D-sphinganine + CO(2).,cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism.,similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.,subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed.,BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Cell Pathway	Endoplasmic reticulum membrane ; Single-pass membrane protein .
CO(2).,cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism.,similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.,subunit:SPTLC1, SPTLC2 and SPTLC3 may encode subunits of the enzyme.,tissue specificity:Widely expressed.,BackgroundThis gene encodes a long chain base subunit of serine palmitoyltransferase. Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Tissue Specificity	Widely expressed.
Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the	Function	CO(2).,cofactor:Pyridoxal phosphate.,pathway:Lipid metabolism; sphingolipid metabolism.,similarity:Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.,subunit:SPTLC1, SPTLC2 and SPTLC3 may encode
	Background	Serine palmitoyltransferase, which consists of two different subunits, is the key enzyme in sphingolipid biosynthesis. It catalyzes the
Nanjing BYabscience technology Co.,Ltd		Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务, 欢迎咨询

	palmitoyl-CoA to 3-oxosphinganine. Mutations in this gene were identified in patients with hereditary sensory neuropathy type I. [provided by RefSeq, Mar 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd