



# EST1 mouse mAb

<b>Catalog No</b>	BYmab-08028
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	CES1 CES2 SES1
<b>Protein Name</b>	EST1
<b>Immunogen</b>	Synthesized peptide derived from human EST1 AA range: 7-57
<b>Specificity</b>	This antibody detects endogenous levels of EST1 at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.142% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Liver carboxylesterase 1 (Acyl-coenzyme A:cholesterol acyltransferase) (ACAT) (Brain carboxylesterase hBr1) (Carboxylesterase 1) (CE-1) (hCE-1) (EC 3.1.1.1) (Cocaine carboxylesterase) (Egasyn) (HMSE) (Methylumbelliferyl-acetate deacetylase 1) (EC 3.1.1.56) (Monocyte/macrophage serine esterase) (Retinyl ester hydrolase) (REH) (Serine esterase 1) (Triacylglycerol hydrolase) (TGH)
<b>Observed Band</b>	60kD
<b>Cell Pathway</b>	Endoplasmic reticulum lumen . Cytoplasm . Lipid droplet . Moves from cytoplasm to lipid droplets upon lipid loading. Associates with lipid droplets independently of triglycerides (TG) content of the droplets and hydrolyzes cholesteryl esters more efficiently from mixed droplets. .
<b>Tissue Specificity</b>	Expressed predominantly in liver with lower levels in heart and lung (PubMed:10562416). Expressed in macrophages (PubMed:11015575, PubMed:21049984, PubMed:18762277).
<b>Function</b>	catalytic activity:A carboxylic ester + H(2)O = an alcohol + a carboxylate.,enzyme regulation:Activated by CHAPS.,function:Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Hydrolyzes

**Nanjing BYabscience technology Co.,Ltd**



aromatic and aliphatic esters, but has no catalytic activity toward amides or a fatty acyl CoA ester.,PTM:Contains sialic acid.,similarity:Belongs to the type-B carboxylesterase/lipase family.,subunit:Homotrimer and homohexamer. Binds to beta-glucuronidase.,tissue specificity:Expressed predominantly in liver with lower levels in heart and lung.,

#### Background

This gene encodes a member of the carboxylesterase large family. The family members are responsible for the hydrolysis or transesterification of various xenobiotics, such as cocaine and heroin, and endogenous substrates with ester, thioester, or amide bonds. They may participate in fatty acyl and cholesterol ester metabolism, and may play a role in the blood-brain barrier system. This enzyme is the major liver enzyme and functions in liver drug clearance. Mutations of this gene cause carboxylesterase 1 deficiency. Three transcript variants encoding three different isoforms have been found for this gene. [provided by RefSeq, Jun 2010],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using EST1 mouse mAb