



EST3 mouse mAb

Catalog No	BYmab-08078
lsotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CES3 UNQ869/PRO1887
Protein Name	EST3
Immunogen	Synthesized peptide derived from human EST3 AA range: 447-497
Specificity	This antibody detects endogenous levels of EST3 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.193% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Carboxylesterase 3 (EC 3.1.1.1) (Liver carboxylesterase 31 homolog)
Observed Band	65kD
Cell Pathway	Endoplasmic reticulum lumen .
Tissue Specificity	Expressed in liver, colon and small intestine.
Function	catalytic activity:A carboxylic ester + H(2)O = an alcohol + a carboxylate.,function:Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Shows low catalytic efficiency for hydrolysis of CPT-11 (7-ethyl-10-[4-(1-piperidino)-1-piperidino]-carbonyloxycamptothecin), a prodrug for camptothecin used in cancer therapeutics.,PTM:N-glycosylated.,similarity:Belongs to the type-B carboxylesterase/lipase family.,tissue specificity:Expressed in liver, colon and small intestine.,
Background	This gene encodes a member of the carboxylesterase large family. The family members are responsible for the hydrolysis or transesterification of various
	Nanjing BYabscience technology Co.,Ltd

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务, 欢迎咨询

	xenobiotics, such as cocaine and heroin, and endogenous substrates with ester, thioester, or amide bonds. They may participate in fatty acyl and cholesterol ester metabolism, and may play a role in the blood-brain barrier system. This gene is expressed in several tissues, particularly in colon, trachea and in brain, and the protein participates in colon and neural drug metabolism. Multiple alternatively spliced transcript variants encoding distinct isoforms have been reported, but the biological validity and/or full-length nature of some variants have not been determined.[provided by RefSeq, Jun 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
Products Images	



Western Blot analysis of various cells using EST3 mouse mAb

Nanjing BYabscience technology Co.,Ltd