



ATS15 mouse mAb

Catalog No	BYmab-08092
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	ADAMTS15
Protein Name	ATS15
Immunogen	Synthesized peptide derived from human ATS15 AA range: 165-215
Specificity	This antibody detects endogenous levels of ATS15 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.207% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	A disintegrin and metalloproteinase with thrombospondin motifs 15 (ADAM-TS 15) (ADAM-TS15) (ADAMTS-15) (EC 3.4.24.-)
Observed Band	105kD
Cell Pathway	Secreted, extracellular space, extracellular matrix . Cell surface .
Tissue Specificity	Expressed in fetal liver and kidney, but not in any of the adult tissues examined.
Function	cofactor: Binds 1 zinc ion per subunit., domain: The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme., domain: The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix., PTM: The precursor is cleaved by a furin endopeptidase., similarity: Contains 1 disintegrin domain., similarity: Contains 1 peptidase M12B domain., similarity: Contains 3 TSP type-1 domains., tissue specificity: Expressed in fetal liver and kidney, but not in any of the adult tissues examined.,

Nanjing BYabscience technology Co.,Ltd



Background

This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs) protein family. ADAMTS family members share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The encoded preproprotein is proteolytically processed to generate the mature enzyme, which may play a role in versican processing during skeletal muscle development. This gene may function as a tumor suppressor in colorectal and breast cancers. [provided by RefSeq, May 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ATS15 mouse mAb