



SEM4G mouse mAb

Catalog No BYmab-08643 Isotype IgG Reactivity Human; Mouse Applications WB Gene Name SEMA4G KIAA1619 Protein Name SEM4G Immunogen Synthesized peptide derived from human SEM4G AA range: 387-437 Specificity This antibody detects endogenous levels of SEM4G at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance, similarity:Belongs to the semaphorin armily, similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain., similarity:Contains 1 Sema domain., similarity:Contains 1 Sema domain., similarity:Contains 1 Sema domain., similarity:Contains 1 Sema domain. Based on sequence and structural similarities, semaphorins are put int		
Reactivity	Catalog No	BYmab-08643
Applications Gene Name SEMA4G KIAA1619 Protein Name SEM4G Immunogen Synthesized peptide derived from human SEM4G AA range: 387-437 Specificity This antibody detects endogenous levels of SEM4G at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function: May play a role in axon guidance _similarity. Belongs to the semaphorin family_similarity. Contains 1 PSI domain_, similarity. Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 1-3. Semaphorins and locations. Multiple transcript receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if not allowed as axon guidance ligands via multimeric receptor complexes, some (if n	Isotype	IgG
Gene Name SEMA4G KIAA1619 Protein Name SEM4G Immunogen Synthesized peptide derived from human SEM4G AA range: 387-437 Specificity This antibody detects endogenous levels of SEM4G at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance., similarity:Belongs to the semaphorin family, similarity:Contains 1 19-like C2-type (immunoglobulin-like) domain., similarity:Contains 1 Tely domain, similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (Sundia plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimerir receptor complexes, some (If not all) containing plexin proteins. Multiple transcript receptor complexes, some (If not all) containing plexin proteins. Multiple transcript receptor complexes, some (If not all) containing plexin proteins. Multiple transcript	Reactivity	Human; Mouse
Protein Name SEM4G Immunogen Synthesized peptide derived from human SEM4G AA range: 387-437 Specificity This antibody detects endogenous levels of SEM4G at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function Function function:May play a role in axon guidance, similarity:Belongs to the semaphorin family, similarity:Contains 1 19-like C2-type (immunoglobulin-like) domain, similarity:Contains 1 19-like occupations, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 1 and 2, viruses h	Applications	WB
Immunogen Synthesized peptide derived from human SEM4G AA range: 387-437 Specificity This antibody detects endogenous levels of SEM4G at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function Function function:May play a role in axon guidancesimilarity:Belongs to the semaphorin family, similarity:Contains 1 1g-like C2-type (immunoglobulin-like) domain., similarity:Contains 1 1g-like C2-type (immunoglobulin-like) domain., similarity:Contains 1 1g-like C2-type (immunoglobulin-like) sema domain., semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via m	Gene Name	SEMA4G KIAA1619
Specificity This antibody detects endogenous levels of SEM4G at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function: May play a role in axon guidance. similarity: Belongs to the semaphorin family, similarity: Contains 1 Ig-like C2-type (immunoglobulin-like) domain., similarity: Contains 1 PSI domain., similarity: Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Protein Name	SEM4G
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance, similarity:Belongs to the semaphorin family, similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain, similarity:Contains 1 Sema domain. Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Immunogen	Synthesized peptide derived from human SEM4G AA range: 387-437
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain. Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3 - Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcriptic letranscriptic perspective complexes, some (if not all) containing plexin proteins. Multiple transcriptic letranscriptic letranscriptic letranscriptic letranscriptic letranscriptic letranscriptic	Specificity	This antibody detects endogenous levels of SEM4G at Human/Mouse
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family,,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain and a PSI domain (found in plexins, semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance, similarity:Belongs to the semaphorin family, similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain., similarity:Contains 1 PSI domain., similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function: May play a role in axon guidance., similarity: Belongs to the semaphorin family, similarity: Contains 1 Ig-like C2-type (immunoglobulin-like) domain., similarity: Contains 1 PSI domain., similarity: Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural simillarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Dilution	WB 1:500-2000
Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Purity	≥90%
Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Single-pass type I membrane protein. Tissue Specificity Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Synonyms	
Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Observed Band	
Function function:May play a role in axon guidance.,similarity:Belongs to the semaphorin family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Cell Pathway	Cell membrane; Single-pass type I membrane protein.
family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 PSI domain.,similarity:Contains 1 Sema domain., Background Semaphorins are a large family of conserved secreted and membrane associated proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Tissue Specificity	
proteins which possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Based on sequence and structural similarities, semaphorins are put into eight classes: invertebrates contain classes 1 and 2, viruses have class V, and vertebrates contain classes 3-7. Semaphorins serve as axon guidance ligands via multimeric receptor complexes, some (if not all) containing plexin proteins. Multiple transcript	Function	family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like)
	Background	

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



RefSeq, Feb 2011],

matters	needing	
attention		

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

