



## FUCO2 mouse mAb

BYmab-08692
Britial 00002
IgG
Human; Mouse;Rat
WB
FUCA2 PSEC0151 UNQ227/PRO260
FUCO2
Synthesized peptide derived from human FUCO2 AA range: 409-459
This antibody detects endogenous levels of FUCO2 at Human/Mouse/Rat
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
Secreted .
catalytic activity:An alpha-L-fucoside + H(2)O = L-fucose + an alcohol.,function:Alpha-L-fucosidase is responsible for hydrolyzing the alpha-1,6-linked fucose joined to the reducing-end N-acetylglucosamine of the carbohydrate moieties of glycoproteins.,similarity:Belongs to the glycosyl hydrolase 29 family.,subunit:Homotetramer.,
This gene encodes a plasma alpha-L-fucosidase, which represents 10-20% of the total cellular fucosidase activity. The protein is a member of the glycosyl hydrolase 29 family, and catalyzes the hydrolysis of the alpha-1,6-linked fucose joined to the reducing-end N-acetylglucosamine of the carbohydrate moieties of glycoproteins. This enzyme is essential for Helicobacter pylori adhesion to human gastric cancer

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using FUCO2 mouse mAb