



# DJC13 mouse mAb

<b>Catalog No</b>	BYmab-08798
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	DNAJC13 KIAA0678 RME8
<b>Protein Name</b>	DJC13
<b>Immunogen</b>	Synthesized peptide derived from human DJC13 AA range: 1576-1626
<b>Specificity</b>	This antibody detects endogenous levels of DJC13 at Human
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Early endosome . Early endosome membrane ; Peripheral membrane protein . Endosome membrane .
<b>Tissue Specificity</b>	
<b>Function</b>	sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 748.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Translated as Glu.,similarity:Contains 1 J domain.,
<b>Background</b>	This gene encodes a member of the Dnaj protein family whose members act as co-chaperones of a partner heat-shock protein by binding to the latter and stimulating ATP hydrolysis. The encoded protein associates with the heat-shock protein Hsc70 and plays a role in clathrin-mediated endocytosis. It may also be involved in post-endocytic transport mechanisms via its associations with other proteins, including the sorting nexin SNX1. Mutations in this gene are associated with Parkinson's disease. [provided by RefSeq, Jun 2016],

**Nanjing BYabscience technology Co.,Ltd**



**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

