



OCAD1 mouse mAb

Catalog No BYmab-08866 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name OCIAD1 OCIA Protein Name OCAD1 Immunogen Synthesized peptide derived from human OCAD1 AA range: 114-164 Specificity This antibody detects endogenous levels of OCAD1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms -20°C/1 year Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity-Belongs to the OCIAD1 family, similarity-Contains 1 OCIA Function similarity-Belongs to the OCIAD1 family, similarity-Contains 1 OCIA		
Reactivity Human; Mouse;Rat Applications WB Gene Name OCIAD1 OCIA Protein Name OCAD1 Immunogen Synthesized peptide derived from human OCAD1 AA range: 114-164 Specificity This antibody detects endogenous levels of OCAD1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monocional, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms - Observed Band Cell Pathway Finsue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland, isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family, similarity:Contains 1 OCIA Function similarity:Belongs to the OCIAD1 family, similarity:Contains 1 OCIA Function similarity:Belongs to the OCIAD1 family, similarity:Contains 1 OCIA <td< td=""><td>Catalog No</td><td>BYmab-08866</td></td<>	Catalog No	BYmab-08866
ApplicationsWBGene NameOCIAD1 OCIAProtein NameOCAD1ImmunogenSynthesized peptide derived from human OCAD1 AA range: 114-164SpecificityThis antibody detects endogenous levels of OCAD1 at Human/Mouse/RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse.IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsEndosome .Cell PathwayEndosome .Tissue SpecificityIsoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord.Functionsimilarity.Eelongs to the OCIAD1 family, similarity.Contains 1 OCIA domain, tissue specificityBackgroundcerebellum and spinal cord.Backgroundsimilarity.eetone to the central nervous system including brain, cerebellum and spinal cord.BackgroundAvoid repeated freezing and thawing!	Isotype	IgG
Gene Name OCIAD1 OCIA Protein Name OCAD1 Immunogen Synthesized peptide derived from human OCAD1 AA range: 114-164 Specificity This antibody detects endogenous levels of OCAD1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Endosome . Tissue Specificity Is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family, similarity:Contains 1 OCIA domain, its uses specificity lissues specificity losform 1 is highly expressed in mamy tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family, similarity:Contains 1 OCIA domain, itsues specifi	Reactivity	Human; Mouse;Rat
Protein Name OCAD1 Immunogen Synthesized peptide derived from human OCAD1 AA range: 114-164 Specificity This antibody detects endogenous levels of OCAD1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Discorm 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 familysimilarity:Contains 1 OCIA domain, issue specificity:Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord., domain, issue specificity:Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord., is attistice to the central nervous system including brain, cerebellum and spinal cord., is restricted to the central nervous system including brain, cerebellum and spinal cord., is restricted to the central nervous system including brain, cerebellum and spinal cord., is restricted	Applications	WB
ImmunogenSynthesized peptide derived from human OCAD1 AA range: 114-164SpecificityThis antibody detects endogenous levels of OCAD1 at Human/Mouse/RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCell PathwayCell PathwayEndosome .Tissue SpecificityIsoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord.Functionsimilarity-Belongs to the OCIAD1 family.similarity:Contains 1 OCIA domain.tissue specificity.Isoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord.,BackgroundAvoid repeated freezing and thawing!	Gene Name	OCIAD1 OCIA
Specificity This antibody detects endogenous levels of OCAD1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family, similarity:Contains 1 OCIA domain, tissue specificity!soform 1 is highly expressed in many tissues, including testis, including testis, is restricted to the central nervous system including brain, cerebellum and spinal cord. Background Mond repeated freezing and thawing!	Protein Name	OCAD1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family., similarity:Contains 1 OCIA domain., tissue specificity:Isoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family., similarity:Contains 1 OCIA domain., tissue specificity:Isoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord. Background cord, matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human OCAD1 AA range: 114-164
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms - Observed Band Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family, similarity:Contains 1 OCIA domain, tissue specificity.Isoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord. Background Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of OCAD1 at Human/Mouse/Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Endosome . Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 familysimilarity:Contains 1 OCIA Background cord Background Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Boogs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity.lsoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord. Background Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord. Background Avoid repeated freezing and thawing!	Purification	5 51 5
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 familysimilarity:Contains 1 OCIA domaintissue specificity:Isoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord. Background Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 familysimilarity:Contains 1 OCIA domaintissue specificity:Isoform 1 is highly expressed in many tissues, including brain, cerebellum and spinal cord. Background Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Background Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Endosome . Cell Pathway Endosome . Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord,, Background Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell PathwayEndosome .Tissue SpecificityIsoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord.Functionsimilarity:Belongs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord,,BackgroundAvoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Function similarity:Belongs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord. Background Avoid repeated freezing and thawing!	Observed Band	
ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord.Functionsimilarity:Belongs to the OCIAD1 family.,similarity:Contains 1 OCIA domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord,.,BackgroundAvoid repeated freezing and thawing!	Cell Pathway	Endosome .
domain.,tissue špecificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal cord, Background matters needing attention	Tissue Specificity	ovary, prostate and mammary gland. Isoform 2 expression is restricted to the
matters needing Avoid repeated freezing and thawing! attention	Function	domain.,tissue specificity:Isoform 1 is highly expressed in many tissues, including testis, brain, placenta, ovary, prostate and mammary gland. Isoform 2 expression is restricted to the central nervous system including brain, cerebellum and spinal
attention	Background	
Nanjing BYabscience technology Co.,Ltd	matters needing attention	Avoid repeated freezing and thawing!
		Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using OCAD1 mouse mAb

Nanjing BYabscience technology Co.,Ltd