



LOXL3 mouse mAb

Catalog No	BYmab-08882
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	LOXL3 LOXL
Protein Name	LOXL3
Immunogen	Synthesized peptide derived from human LOXL3 AA range: 130-180
Specificity	This antibody detects endogenous levels of LOXL3 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Secreted, extracellular space . Cytoplasm . Nucleus . It is unclear how LOXL3 is both intracellular (cytoplasmic and nuclear) and extracellular: it contains a clear signal sequence and is predicted to localize in the extracellular medium. However, the intracellular location is clearly reported and at least another protein of the family (LOXL2) also has intracellular and extracellular localization despite the presence of a signal sequence (PubMed:28065600). .; [Isoform 1]: Secreted, extracellular space .; [Isoform 2]: Cytoplasm . Secreted, extracellular space .
Tissue Specificity	Isoform 1: Predominantly detected in the heart, placenta, lung, and small intestine (PubMed:17018530). Isoform 2: Highly detected in the kidney, pancreas, spleen, and thymus, and is absent in lung (PubMed:17018530). In eye, present in all layers of corneas as well as in the limbus and conjunctiva (at protein level) (PubMed:26218558).
Function	cofactor:Contains 1 lysine tyrosylquinone.,cofactor:Copper.,PTM:The lysine tyrosylquinone cross-link (LTQ) is generated by condensation of the epsilon-amino group of a lysine with a topaquinone produced by oxidation of

Nanjing BYabscience technology Co.,Ltd



tyrosine.,similarity:Belongs to the lysyl oxidase family.,similarity:Contains 4 SRCR domains.,tissue specificity:Expressed in many tissues, the highest levels among the tissues studied being seen in the placenta, heart, ovary, testis, small intestine and spleen.,

Background

This gene encodes a lysyl oxidase, which likely functions as an amine oxidase and plays a role in the formation of crosslinks in collagens and elastin. Deletion of the related gene in mouse causes neonatal mortality with cleft palate, spine deformity, and defects in collagen organization. A mutation in this gene was found in a family with Stickler syndrome. [provided by RefSeq, Sep 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using LOXL3 mouse mAb