



## SPRE3 mouse mAb

Catalog NoBYmab-08917IsotypeIgGReactivityHuman; MouseApplicationsWBGene NameSPRED3 EVE-3Protein NameSPRE33ImmunogenSynthesized peptide derived from human SPRE3 AA range: 311-361SpecificityThis antibody detects endogenous levels of SPRE3 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionW B 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearCherrer BandomCell membrane ; Peripheral membrane protein .Tissue SpecificityUnction: Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase., PTM:Phosphorylated on tyrosine.similarity:Contains 1 SPR (sprouty) domain, similarity:Contains 1 WH1 domain, subutilInteracts with Ras, BackgroundThis gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilar-stimulated protein (MAP) kinase signaling, particularly during organgeneens.Terminal Sprouty-like rysteine-rich domain signaling, particularly during organgeneens.Memory and an N-terminal Sprouty-like rysteine-rich domain signaling, particularly during organgeneens. J Memory and an N-terminal Sprouty-like rysteine-rich domain signaling, particularly during organgeneens. J Memory and an N-terminal Sprouty-like rysteine-rich domain signaling, particularly during organgeneens. J Memory and an N-terminal Sprout		
Reactivity   Human; Mouse     Applications   WB     Gene Name   SPRED3 EVE-3     Protein Name   SPRE3     Immunogen   Synthesized peptide derived from human SPRE3 AA range: 311-361     Specificity   This antibody detects endogenous levels of SPRE3 at Human/Mouse     Formulation   Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.     Source   Monoclonal, Mouse, IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cell membrane ; Peripheral membrane protein .     Tissue Specificity   Function     Function   function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase. JPTM:Phosphorylated on tyrosine. similarity.Contains 1 SPR (SPT) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (MAP) kinase.     Background   This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (MAP) kinase signaling, particularly during organogenesis. A Humani, sublative of a family of proteins in a methode solicin in subtrate solicin in subtrate solicin in subtrate solicin in subtrate solicin muticate solicin in subtrate solicin in subtrate solicin in subtrate solicin genesits in muticate </td <td>Catalog No</td> <td>BYmab-08917</td>	Catalog No	BYmab-08917
Applications       WB         Gene Name       SPRED3 EVE-3         Protein Name       SPRE3         Immunogen       Synthesized peptide derived from human SPRE3 AA range: 311-361         Specificity       This antibody detects endogenous levels of SPRE3 at Human/Mouse         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Cell membrane ; Peripheral membrane protein .         Tissue Specificity       Cell membrane ; Peripheral membrane protein .         Tissue Specificity       function: Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase, PTM.Phosphorylated on tyrosine, similarity.Contains 1 SPR (sprouty) domain, similarity.Contains 1 SPR (sprouty) domain, similarity.Contains 1 SPR (sprouty) -4(EVH-1) domain. The encoded protein is newth factor-mediated activation of MAP kinase. PTM.Phosphorylated on tyrosine, similarity.Contains 1 SPR (sprouty) -4(EVH-1) domain. The encoded protein is newth factor-mediated activation of MAP kinase. PTM.Phosphorylated on tyrosine, simulated protein (MASP) homology-1 (EVH-1) domain. T	lsotype	lgG
Gene Name       SPRED3 EVE-3         Protein Name       SPRE3         Immunogen       Synthesized peptide derived from human SPRE3 AA range: 311-361         Specificity       This antibody detects endogenous levels of SPRE3 at Human/Mouse         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Cell membrane ; Peripheral membrane protein .         Tissue Specificity       Function: Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase, PTM:Phosphorylated on tyrosine, similarity:Contains 1 SPR (SPY) and an N-terminal Ena/Vasodilator-stimulated phosphorptoein (VASP) homology (YASP) homonology (YASP) homology (YASP	Reactivity	Human; Mouse
Protein Name       SPRE3         Immunogen       Synthesized peptide derived from human SPRE3 AA range: 311-361         Specificity       This antibody detects endogenous levels of SPRE3 at Human/Mouse         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -         Cell Pathway       Cell membrane ; Peripheral membrane protein .         Tissue Specificity       function: Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase, PTM:Phosphorylated on tyrosine, similarity:Contains 1 SPR (sproty) domain "similarity.Contains 1 SPR (sproty) domain "ginalitarty-contains 1 MRAs, Background         This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing result in multiple	Applications	WB
ImmunogenSynthesized peptide derived from human SPRE3 AA range: 311-361SpecificityThis antibody detects endogenous levels of SPRE3 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCell membrane ; Peripheral membrane protein .Tissue SpecificityCell membrane ; Peripheral membrane protein .Functionfunction:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 MVH domain. subunit:Interacts with Ras., BackgroundBackgroundThis gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodliator-studied photeping (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (VASP) kinase signaling, particulaty during organogeneasis. Alternative splicing results in multiple	Gene Name	SPRED3 EVE-3
Specificity     This antibody detects endogenous levels of SPRE3 at Human/Mouse       Formulation     Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.       Source     Monoclonal, Mouse,IgG       Purification     The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.       Dilution     WB 1:500-2000       Concentration     1 mg/ml       Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     Cell membrane ; Peripheral membrane protein .       Function     function: Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase., PTM:Phosphorylated on tyrosine, similarity:Contains 1 SPR (sprouty) domain., similarity:Contains 1 WH1 domain., subunit.Interacts with Ras., Background       This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain SRY) and an N-terminal Ena/Vasciliator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Protein Name	SPRE3
Formulation     Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.       Source     Monoclonal, Mouse,IgG       Purification     The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.       Dilution     WB 1:500-2000       Concentration     1 mg/ml       Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     -20°C/1 year       Observed Band     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     Function: Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase, PTM:Phosphorylated on tyrosine, similarity:Contains 1 SPR (sprouty) domain., similarity:Contains 1 SPR (Sprouty) domain., similarity:Contains 1 WH1 domain., subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphorptoein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase singing, particularly during organogenesis. Alternative splicing results in multiple	Immunogen	Synthesized peptide derived from human SPRE3 AA range: 311-361
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Cell Pathway         Cell Pathway       Cell membrane ; Peripheral membrane protein .         Tissue Specificity       function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.PTM:Phosphorylated on tyrosine, similarity:Contains 1 SPR (sprouty) domain., similarity:Contains 1 WH1 domain, subunit:Interacts with Ras.,         Background       This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (MAP) kinase, proteins 1 WH1 domain, subunit:Interacts with Ras.,	Specificity	This antibody detects endogenous levels of SPRE3 at Human/Mouse
Purification     The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.       Dilution     WB 1:500-2000       Concentration     1 mg/ml       Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     -20°C/1 year       Observed Band     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     Function of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit.Interacts with Ras., Background       This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain the encoded protein (MAP) kinase., Proteins that negatively regulates mitogen-activated protein (MAP) kinase substrate membrane.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.       Dilution     WB 1:500-2000       Concentration     1 mg/ml       Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     -20°C/1 year       Observed Band     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     Function       Function     function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase., PTM:Phosphorylated on tyrosine., similarity:Contains 1 SPR (sprouty) domain., similarity:Contains 1 WH1 domain., subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively using during organogenesis. Alternative splicing results in multiple	Source	Monoclonal, Mouse,IgG
Concentration     1 mg/ml       Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     -20°C/1 year       Observed Band     -20°C/1 year       Cell Pathway     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     -20°C/1 year       Function     function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Purification	
Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     -20°C/1 year       Observed Band     -20°C/1 year       Cell Pathway     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     -20°C/1 year       Function     function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Dilution	WB 1:500-2000
Storage Stability     -20°C/1 year       Synonyms     -20°C/1 year       Observed Band     -20°C/1 year       Cell Pathway     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     -20°C/1 year       Function     function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Concentration	1 mg/ml
Synonyms       Observed Band       Cell Pathway     Cell membrane ; Peripheral membrane protein .       Tissue Specificity       Function     function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Purity	≥90%
Observed Band       Cell Pathway     Cell membrane ; Peripheral membrane protein .       Tissue Specificity       Function     function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Storage Stability	-20°C/1 year
Cell Pathway     Cell membrane ; Peripheral membrane protein .       Tissue Specificity     Tissue Specificity       Function     function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,       Background     This gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Synonyms	
Tissue SpecificityFunctionfunction:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,BackgroundThis gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Observed Band	
Functionfunction:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,BackgroundThis gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Cell Pathway	Cell membrane ; Peripheral membrane protein .
of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,BackgroundThis gene encodes a protein with a C-terminal Sprouty-like cysteine-rich domain (SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Tissue Specificity	
(SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple	Function	function:Tyrosine kinase substrate that inhibits growth-factor-mediated activation of MAP kinase.,PTM:Phosphorylated on tyrosine.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subunit:Interacts with Ras.,
	Background	(SRY) and an N-terminal Ena/Vasodilator-stimulated phosphoprotein (VASP) homology-1 (EVH-1) domain. The encoded protein is a member of a family of proteins that negatively regulates mitogen-activated protein (MAP) kinase signaling, particularly during organogenesis. Alternative splicing results in multiple

Nanjing BYabscience technology Co.,Ltd






## **matters needing** Avoid repeated freezing and thawing! attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Nanjing BYabscience technology Co.,Ltd