



# KIF1B mouse mAb

<b>Catalog No</b>	BYmab-09033
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	KIF1B KIAA0591 KIAA1448
<b>Protein Name</b>	KIF1B
<b>Immunogen</b>	Synthesized peptide derived from human KIF1B AA range: 1331-1381
<b>Specificity</b>	This antibody detects endogenous levels of KIF1B at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton. Mitochondrion . Cell projection, axon .; [Isoform 1]: Cytoplasmic vesicle, secretory vesicle, synaptic vesicle .
<b>Tissue Specificity</b>	Isoform 3 is abundant in the skeletal muscle. It is also expressed in fetal brain, lung and kidney, and adult heart, placenta, testis, ovary and small intestine. Isoform 2 is abundant in the brain and also expressed in fetal heart, lung, liver and kidney, and adult skeletal muscle, placenta, liver, kidney, heart, spleen, thymus, prostate, testis, ovary, small intestine, colon and pancreas.
<b>Function</b>	disease:Defects in KIF1B are the cause of Charcot-Marie-Tooth disease type 2A1 (CMT2A1) [MIM:118210]. CMT2A1 is a form of Charcot-Marie-Tooth disease, the most common inherited disorder of the peripheral nervous system. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathy or CMT1, and primary peripheral axonal neuropathy or CMT2. Neuropathies of the CMT2 group are characterized by signs of axonal regeneration in the absence of obvious myelin alterations, normal or slightly

**Nanjing BYabscience technology Co.,Ltd**



reduced nerve conduction velocities, and progressive distal muscle weakness and atrophy.,function:Motor for anterograde transport of mitochondria. Has a microtubule plus end-directed motility.,similarity:Belongs to the kinesin-like protein family.,similarity:Belongs to the kinesin-like protein family. U

#### Background

This gene encodes a motor protein that transports mitochondria and synaptic vesicle precursors. Mutations in this gene cause Charcot-Marie-Tooth disease, type 2A1. [provided by RefSeq, Jul 2008],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using KIF1B mouse mAb