



MAGI3 mouse mAb

36
ise;Rat
1634
peptide derived from human MAGI3 AA range: 927-977
y detects endogenous levels of MAGI3 at Human/Mouse/Rat
S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Mouse,IgG
v was affinity-purified from mouse antiserum by natography using epitope-specific immunogen.
000
ne; Peripheral membrane protein. Cell junction, tight junction. Incentrates in specific sites at the plasma membrane and in the pithelial cells, it localizes at tight junctions (By similarity)
essed.
as a scaffolding protein at cell-cell junctions, thereby regulating lar and signaling processes. Cooperates with PTEN to modulate the ty of AKT1. Its interaction with PTPRB and tyrosine phosphorylated gests that it may link receptor tyrosine phosphatase with its the plasma membrane. In polarized epithelial cells, involved in cking of TGFA to the cell surface. Regulates the ability of LPAR2 to and RhoA pathways. Regulates the JNK signaling cascade via its ith FZD4 and VANGL2.,miscellaneous:MAGI3 PDZ domains are gn peptide ligands that bind and inhibit PDZ M:Ubiquitinated following interaction with HPV E6 protein, leading to on by the proteasome. Degradation is independent of E6AP ubiquitin

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





ligase complex., sequence caution: Translation N-terminally

Background	
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Nanjing BYabscience technology Co.,Ltd