



PARG mouse mAb

Catalog No	BYmab-09106
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	PARG
Protein Name	PARG
Immunogen	Synthesized peptide derived from human PARG AA range: 140-190
Specificity	This antibody detects endogenous levels of PARG at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year [Isoform 1]: Nucleus . Colocalizes with PCNA at replication foci (PubMed:21398629). Relocalizes to the cytoplasm in response to DNA damage (PubMed:16460818); [Isoform 2]: Cytoplasm . Translocates to the nucleus in response to DNA damage; [Isoform 3]: Cytoplasm .; [Isoform 4]: Cytoplasm .
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year [Isoform 1]: Nucleus . Colocalizes with PCNA at replication foci (PubMed:21398629). Relocalizes to the cytoplasm in response to DNA damage (PubMed:16460818); [Isoform 2]: Cytoplasm . Translocates to the nucleus in response to DNA damage; [Isoform 3]: Cytoplasm .; [Isoform 4]: Cytoplasm . Mitochondrion .; [Isoform 5]: Mitochondrion matrix .

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	activation of the antiviral prodrug valacyclovir. Alternatively spliced transcript variants have been described.[provided by RefSeq, Jan 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



