




DBR1 mouse mAb

BYmab-09125
IgG
Human; Mouse
WB
DBR1
DBR1
Synthesized peptide derived from human DBR1 AA range: 230-280
This antibody detects endogenous levels of DBR1 at Human/Mouse
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
Nucleus .
Ubiquitously expressed, strongest expression in the spinal cord and brainstem.
catalytic activity:RNA processing activity that hydrolyzes the 2'-5' phosphodiester linkage at the branchpoint of excised intron lariats.,cofactor:Divalent metal cations.,function:Cleaves the 2'-5' phosphodiester linkage at the branch point of lariat intron pre-mRNAs after splicing and converts them into linear molecules that are subsequently degraded. It thereby facilitates ribonucleotide turnover. It may also participate in retrovirus replication via a RNA lariat intermediate in cDNA synthesis.,miscellaneous:Cells lacking DBR1 show an inhibition of HIV-1 replication.,similarity:Belongs to the lariat debranching enzyme family.,
The protein encoded by this gene is an RNA lariat debranching enzyme that hydrolyzes 2'-5' prime branched phosphodiester bonds. The encoded protein specifically targets the bonds at the branch point of excised lariat

Nanjing BYabscience technology Co.,Ltd







	intron RNA, converting them to linear molecules that are then degraded. This protein may also be involved in retroviral replication. [provided by RefSeq, Nov 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

