



# ZC12D mouse mAb

Catalog No	BYmab-09274
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	ZC3H12D C6orf95 MCPIP4 TFL
Protein Name	Probable ribonuclease ZC3H12D (EC 3.1.-.-) (MCP-induced protein 4) (Transformed follicular lymphoma) (Zinc finger CCCH domain-containing protein 12D) (p34)
Immunogen	Synthesized peptide derived from human ZC12D AA range: 87-137
Specificity	This antibody detects endogenous levels of human ZC12D
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm, P-body . Colocalizes with ZC3H12A in GW bodies (GWBs) (PubMed:26134560). .; [Isoform 1]: Cytoplasm. Localized as discrete granules. Colocalized with mRNA-processing body markers, AGO2 and DCP1A, but not with a stress granule maker, TIA1, in the cytoplasm.; [Isoform 3]: Cytoplasm. Nucleus.
Tissue Specificity	Expressed in normal human lymphocytes but defective in some leukemia/lymphoma cell lines.
Function	
Background	

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658



**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images