



IFT20 mouse mAb

Catalog No	BYmab-09458
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	IFT20
Protein Name	Intraflagellar transport protein 20 homolog (hIFT20)
Immunogen	Synthesized peptide derived from human IFT20 AA range: 62-112
Specificity	This antibody detects endogenous levels of human IFT20
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	spermatids. In step 14 spermatids, found in the basal body of the sperm tail.
Cell Pathway Tissue Specificity	organizing center, centrosome, centriole . Cytoplasm, cytoskeleton, cilium basal body . Cell projection, cilium . Cytoplasm, cytoskeleton . Golgi apparatus . Cytoplasmic vesicle, secretory vesicle, acrosome . Cytoplasm . Present at the centrosomes during the cell cycle and associated with the proximal portion of the mother centriole and the lateral aspect of the daughter centriole. Associated with basal body at the base of primary cilia. Detected in the Golgi apparatus of round spermatids and late spermatocytes. Also detected in the manchette of step 10-12 spermatids. In step 14 spermatids, found in the basal body of the sperm tail.
	organizing center, centrosome, centriole . Cytoplasm, cytoskeleton, cilium basal body . Cell projection, cilium . Cytoplasm, cytoskeleton . Golgi apparatus . Cytoplasmic vesicle, secretory vesicle, acrosome . Cytoplasm . Present at the centrosomes during the cell cycle and associated with the proximal portion of the mother centriole and the lateral aspect of the daughter centriole. Associated with basal body at the base of primary cilia. Detected in the Golgi apparatus of round spermatids and late spermatocytes. Also detected in the manchette of step 10-12 spermatids. In step 14 spermatids, found in the basal body of the sperm tail. Localization in the manchette of elongating spermatids is dependent on SPAG17.

Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998





Background

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
Products Images	

Nanjing BYabscience technology Co.,Ltd