



ROGDI mouse mAb

Catalog No	BYmab-09643
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ROGDI
Protein Name	Protein rogdi homolog
Immunogen	Synthesized peptide derived from human ROGDI AA range: 148-198
Specificity	This antibody detects endogenous levels of human ROGDI
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Dilution Concentration	WB 1:500-2000 1 mg/ml
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Concentration Purity Storage Stability	1 mg/ml ≥90%
Concentration Purity Storage Stability Synonyms	1 mg/ml ≥90% -20°C/1 year Nucleus envelope . Cell junction, synapse, presynapse . Cell projection, axon . Perikaryon . Cell projection, dendrite . Cytoplasmic vesicle, secretory vesicle,
Concentration Purity Storage Stability Synonyms Observed Band	1 mg/ml ≥90% -20°C/1 year Nucleus envelope . Cell junction, synapse, presynapse . Cell projection, axon . Perikaryon . Cell projection, dendrite . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . Detected primarily at presynaptic sites on axons, and to a lesser
Concentration Purity Storage Stability Synonyms Observed Band Cell Pathway	1 mg/ml ≥90% -20°C/1 year Nucleus envelope . Cell junction, synapse, presynapse . Cell projection, axon . Perikaryon . Cell projection, dendrite . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . Detected primarily at presynaptic sites on axons, and to a lesser degree in soma and dendrites. Not detected at post-synaptic sites Widely expressed with highest levels in spinal cord, brain, heart and bone
Concentration Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	1 mg/ml ≥90% -20°C/1 year Nucleus envelope . Cell junction, synapse, presynapse . Cell projection, axon . Perikaryon . Cell projection, dendrite . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . Detected primarily at presynaptic sites on axons, and to a lesser degree in soma and dendrites. Not detected at post-synaptic sites Widely expressed with highest levels in spinal cord, brain, heart and bone

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd