



PARVG mouse mAb

Catalog No	BYmab-09869
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	PARVG
Protein Name	Gamma-parvin
Immunogen	Synthesized peptide derived from human PARVG AA range: 113-163
Specificity	This antibody detects endogenous levels of human PARVG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cell junction, focal adhesion. Cell membrane ; Peripheral membrane protein ;
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Constituent of focal adhesions. Expressed predominantly in lymphoid organs, including spleen, thymus, lymph node, bone marrow and peripheral blood leukocytes and moderately in the digestive tract, including stomach, duodenum, jejunum, ileum, ileocecum and appendix, as well as in lung and liver. Also expressed in tumors, but at a lower
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Constituent of focal adhesions. Expressed predominantly in lymphoid organs, including spleen, thymus, lymph node, bone marrow and peripheral blood leukocytes and moderately in the digestive tract, including stomach, duodenum, jejunum, ileum, ileocecum and appendix, as well as in lung and liver. Also expressed in tumors, but at a lower
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Constituent of focal adhesions. Expressed predominantly in lymphoid organs, including spleen, thymus, lymph node, bone marrow and peripheral blood leukocytes and moderately in the digestive tract, including stomach, duodenum, jejunum, ileum, ileocecum and appendix, as well as in lung and liver. Also expressed in tumors, but at a lower

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd