



Bmi-1 mouse Monoclonal Antibody(F270)

Catalog No	BYmab-10287
Isotype	IgG
Reactivity	Human
Applications	WB
Gene Name	BMI1 PCGF4 RNF51
Protein Name	Bmi-1
Immunogen	Recombinant Protein derived from human Bmi-1
Specificity	This antibody detects endogenous levels of human Bmi-1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Polycomb complex protein BMI-1 (Polycomb group RING finger protein 4) (RING finger protein 51)
Observed Band	37-43kD
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Epithelium,Erythrocyte,Muscle,Thymus,
Function	disease:Cooperates with the MYC oncogene to produce B-lymphomas.,function:Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. In the PRC1 complex, it is required to stimulate the E3 ubiquitin-protein ligase activity of RNF2/RING2.,PTM:May be polyubiquitinated; which does not lead to proteasomal degradation.,similarity:Contains 1 RING-type zinc finger.,subunit:Component of chromatin-associated class II polycomb repressive complex 1 (PRC1/hPRC-H) at

Nanjing BYabscience technology Co.,Ltd



least composed of PCGF2/RNF110, BMI1/PCGF4, CBX2/M33, CBX4/PC2, CBX8/PC3, PHC1, PHC2, PHC3, SCMH1, RING1 and R

Background

BMI1 proto-oncogene, polycomb ring finger(BMI1) Homo sapiens This gene encodes a ring finger protein that is major component of the polycomb group complex 1 (PRC1). This complex functions through chromatin remodeling as an essential epigenetic repressor of multiple regulatory genes involved in embryonic development and self-renewal in somatic stem cells. This protein also plays a central role in DNA damage repair. This gene is an oncogene and aberrant expression is associated with numerous cancers and is associated with resistance to certain chemotherapies. A pseudogene of this gene is found on chromosome X. Read-through transcription also exists between this gene and the upstream COMM domain containing 3 (COMMD3) gene. [provided by RefSeq, Sep 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658